

DEPARTMENT OF COMMERCE**National Oceanic and Atmospheric Administration**

RIN 0648–XD727

Takes of Marine Mammals Incidental to Specified Activities; Low-Energy Marine Geophysical Survey in the Southwest Pacific Ocean, East of New Zealand, May to June 2015

AGENCY: National Marine Fisheries Service (NMFS), National Oceanic and Atmospheric Administration (NOAA), Commerce.

ACTION: Notice; proposed Incidental Harassment Authorization; request for comments.

SUMMARY: NMFS has received an application from the Scripps Institution of Oceanography (SIO), on behalf of SIO and the U.S. National Science Foundation (NSF), for an Incidental Harassment Authorization (IHA) to take marine mammals, by harassment, incidental to conducting a low-energy marine geophysical (seismic) survey in the Southwest Pacific Ocean, East of New Zealand, May to June 2015. Pursuant to the Marine Mammal Protection Act (MMPA), NMFS is requesting comments on its proposal to issue an IHA to SIO to incidentally harass, by Level B harassment only, 32 species of marine mammals during the specified activity.

DATES: Comments and information must be received no later than April 20, 2015.

ADDRESSES: Comments on the application should be addressed to Jolie Harrison, Chief, Permits and Conservation Division, Office of Protected Resources, National Marine Fisheries Service, 1315 East-West Highway, Silver Spring, MD 20910. The mailbox address for providing email comments is ITP.Goldstein@noaa.gov. Please include 0648–XD727 in the subject line. NMFS is not responsible for email comments sent to addresses other than the one provided here. Comments sent via email, including all attachments, must not exceed a 25-megabyte file size.

Instructions: All comments received are a part of the public record and will generally be posted to: <http://www.nmfs.noaa.gov/pr/permits/incidental/> without change. All Personal Identifying Information (for example, name, address, etc.) voluntarily submitted by the commenter may be publicly accessible. Do not submit Confidential Business Information or otherwise sensitive or protected information.

A copy of the IHA application may be obtained by writing to the address specified above, telephoning the contact listed here (see **FOR FURTHER INFORMATION CONTACT**) or visiting the Internet at: <http://www.nmfs.noaa.gov/pr/permits/incidental/>. Documents cited in this notice may also be viewed by appointment, during regular business hours, at the aforementioned address.

A “Draft Environmental Analysis of a Low-Energy Marine Geophysical Survey by the R/V *Roger Revelle* in the Southwest Pacific Ocean, East of New Zealand, May to June 2015” (Draft Environmental Analysis) in accordance with the National Environmental Policy Act (NEPA) and the regulations published by the Council of Environmental Quality (CEQ), has been prepared on behalf of NSF and SIO. It is posted at the foregoing site. NMFS has independently evaluated the Draft Environmental Analysis and has prepared a separate NEPA analysis titled “Draft Environmental Assessment on the Issuance of an Incidental Harassment Authorization to the Scripps Institution of Oceanography to Take Marine Mammals by Harassment Incidental to a Low-Energy Marine Geophysical Survey in the Southwest Pacific Ocean, East of New Zealand, May to June 2015.” Information in the SIO’s IHA application, Draft Environmental Analysis, Draft EA and this notice of the proposed IHA collectively provide the environmental information related to proposed issuance of the IHA for public review and comment. NMFS will review all comments submitted in response to this notice as we complete the NEPA process, including a decision of whether to sign a Finding of No Significant Impact (FONSI), prior to a final decision on the IHA request.

FOR FURTHER INFORMATION CONTACT: Howard Goldstein or Jolie Harrison, Office of Protected Resources, NMFS, 301–427–8401.

SUPPLEMENTARY INFORMATION:**Background**

Sections 101(a)(5)(A) and (D) of the MMPA, (16 U.S.C. 1361 *et seq.*) direct the Secretary of Commerce (Secretary) to allow, upon request, the incidental, but not intentional, taking of small numbers of marine mammals by United States citizens who engage in a specified activity (other than commercial fishing) within a specified geographical region if certain findings are made and either regulations are issued or, if the taking is limited to harassment, a notice of a proposed authorization is provided to the public for review.

An authorization for incidental takings shall be granted if NMFS finds that the taking will have a negligible impact on the species or stock(s), will not have an unmitigable adverse impact on the availability of the species or stock(s) for subsistence uses (where relevant), and if the permissible methods of taking and requirements pertaining to the mitigation, monitoring and reporting of such takings are set forth. NMFS has defined “negligible impact” in 50 CFR 216.103 as “. . . an impact resulting from the specified activity that cannot be reasonably expected to, and is not reasonably likely to, adversely affect the species or stock through effects on annual rates of recruitment or survival.”

Section 101(a)(5)(D) of the MMPA established an expedited process by which citizens of the United States can apply for an authorization to incidentally take small numbers of marine mammals by harassment. Section 101(a)(5)(D) of the MMPA establishes a 45-day time limit for NMFS’s review of an application, followed by a 30-day public notice and comment period on any proposed authorizations for the incidental harassment of small numbers of marine mammals. Within 45 days of the close of the public comment period, NMFS must either issue or deny the authorization.

Except with respect to certain activities not pertinent here, the MMPA defines “harassment” as: any act of pursuit, torment, or annoyance which (i) has the potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment]; or (ii) has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering [Level B harassment].

Summary of Request

On December 15, 2014, NMFS received an application from SIO, on behalf of SIO and NSF, requesting that NMFS issue an IHA for the take, by Level B harassment only, of small numbers of marine mammals incidental to conducting a low-energy marine seismic survey as well as heat-flow measurements in the Southwest Pacific Ocean, at three sites off the east coast of New Zealand, during May to June 2015. The sediment coring component of the proposed project, which was described in the IHA application and Draft Environmental Analysis, was not funded and no piston or gravity coring for seafloor samples would be

conducted during the low-energy seismic survey. The low-energy seismic survey would take place within the Exclusive Economic Zone (EEZ) and outside the territorial waters of New Zealand. On behalf of SIO, the U.S. Department of State is seeking authorization from New Zealand for clearance to work within the EEZ.

The research would be conducted by Oregon State University and funded by the U.S. National Science Foundation (NSF). SIO plan to use one source vessel, the R/V *Roger Revelle* (*Revelle*), and a seismic airgun array and hydrophone streamer to collect seismic data in the Southwest Pacific Ocean, East of New Zealand. SIO plans to use conventional low-energy, seismic methodology to perform marine-based studies in the Southwest Pacific Ocean (see Figure 1 of the IHA application). The studies would involve a low-energy seismic survey and heat-flow measurements from the seafloor to meet a number of research goals. In addition to the proposed operations of the seismic airgun array and hydrophone streamer, SIO intends to operate two additional acoustical data acquisition systems—a multi-beam echosounder and sub-bottom profiler continuously throughout the low-energy seismic survey.

Acoustic stimuli (*i.e.*, increased underwater sound) generated during the operation of the seismic airgun array have the potential to cause behavioral disturbance for marine mammals in the proposed study area. This is the principal means of marine mammal taking associated with these activities, and SIO have requested an authorization to take 32 species of marine mammals by Level B harassment. Take is not expected to result from the use of the multi-beam echosounder and sub-bottom profiler, as the brief exposure of marine mammals to one pulse, or small numbers of signals, to be generated by these

instruments in this particular case is not likely to result in the harassment of marine mammals. Also, NMFS does not expect take to result from collision with the source vessel because it is a single vessel moving at a relatively slow, constant cruise speed of 5 knots ([kts]; 9.3 kilometers per hour [km/hr]; 5.8 miles per hour [mph]) during seismic acquisition within the study area, for a relatively short period of time (approximately 27 operational days). It is likely that any marine mammal would be able to avoid the vessel.

Description of the Proposed Specified Activity

Overview

SIO proposes to use one source vessel, the *Revelle*, a two GI airgun array and one hydrophone streamer to conduct the conventional seismic survey as part of the NSF-funded research project “Collaborative Research: The Thermal Regime of the Hikurangi Subduction Zone and Shallow Slow Slip Events, New Zealand.” In addition to the airguns, SIO intends to conduct a bathymetric survey and heat-flow measurements at three sites off the southwest coast of North Island and northeast coast of South Island, New Zealand from the *Revelle* during the proposed low-energy seismic survey.

Proposed Dates and Duration

The *Revelle* is expected to depart from Auckland, New Zealand on approximately May 18, 2015 and arrive at Napier, New Zealand on approximately June 18, 2015. Airgun operations would take approximately 135 hours in total, and the remainder of the time would be spent in transit and collecting heat-flow measurements and cores. The total distance the *Revelle* would travel in the region to conduct the proposed research activities (*i.e.*, seismic survey, bathymetric survey, and transit to heat-flow measurement locations) represents approximately


2,000 km (1,079.9 nmi). Some minor deviation from this schedule is possible, depending on logistics and weather (*e.g.*, the cruise may depart earlier or be extended due to poor weather; or there could be additional days of airgun operations if collected data are deemed to be of substandard quality).

Proposed Specified Geographic Region

The proposed project and survey sites are located off the southeast coast of North Island and northeast coast of the South Island, New Zealand in selected regions of the Southwest Pacific Ocean. The proposed survey sites are located between approximately 38.5 to 42.5° South and approximately 174 to 180° East off the east coast of New Zealand, in the EEZ of New Zealand and outside of territorial waters (see Figure 1 of the IHA Application). Water depths in the study area are between approximately 200 to 3,000 m (656.2 to 9,842.5 ft). The proposed low-energy seismic survey would be collected in a total of nine grids of intersecting lines of two sizes (see Figure 1 of the IHA application) at exact locations to be determined in the field during May to June 2015. Figure 1 also illustrates the general bathymetry of the proposed study area. The proposed low-energy seismic survey would be within an area of approximately 1,154 km² (336.5 nmi²). This estimate is based on the maximum number of kilometers for the low-energy seismic survey (1,250 km) multiplied by the area ensonified around the planned tracklines (2 x 0.6 km in intermediate water depths and 2 x 0.4 km in deep water depths). The ensonified area is based on the predicted rms radii (m) based on modeling and empirical measurements (assuming 100% use of the two 45 in³ GI airguns in 100 to 1,000 m or greater than 1,000 m water depths), which was calculated to be 600 m (1,968.5 ft) or 400 m (1,312.3 ft).

BILLING CODE 3510-22-P

Figure 1. Locations of the proposed low-energy seismic survey and heat-flow probe measurement sites east of New Zealand, May to June 2015.



BILLING CODE 3510-22-C

Detailed Description of the Proposed Specified Activity

In support of a research project put forward by Oregon State University (OSU) and to be funded by NSF, SIO proposes to conduct a low-energy seismic survey in the Southwest Pacific

Ocean, East of New Zealand, from May to June 2015. In addition to the low-energy seismic survey, scientific research activities would include conducting a bathymetric profile survey of the seafloor using transducer-based instruments such as a multi-beam echosounder and sub-bottom profiler;

and heat-flow measurements from the seafloor using various methods and equipment at three sites off the southeast coast of North Island and northeast coast of South Island, New Zealand. Water depths in the survey area are approximately 200 to 3,000 meters (m) (656.2 to 9,842.5 feet [ft]).

The proposed low-energy seismic survey is scheduled to occur for a total of approximately 135 hours over the course of the entire cruise, which would be for approximately 27 operational days in May to June 2015. The proposed low-energy seismic survey would be conducted during the day (from nautical twilight-dawn to nautical twilight-dusk) and night, and for up to approximately 72 hours of continuous operations at a time. The operation hours and survey length would include equipment testing, ramp-up, line changes, and repeat coverage. Some minor deviation from these dates would be possible, depending on logistics and weather. The Principal Investigators are Dr. R. N. Harris and Dr. A. Trehu of the OSU.

The proposed surveys would allow the development of a process-based understanding of the thermal structure of the Hikurangi subduction zone, and the expansion of this understanding by using regional observations of gas hydrate-related bottom simulating reflections. To achieve the proposed project's goals, the Principal Investigators propose to collect low-energy, high-resolution multi-channel system profiles, heat-flow measurements, and sediment cores along transects seaward and landward of the Hikurangi deformation front. Heat-flow measurements would be made in well-characterized sites, increasing the number of publicly available heat-flow and thermal conductivity measurements from this continental margin by two orders of magnitude. Seismic survey data would be used to produce sediment structural maps and seismic velocities to achieve the project objectives. Data from sediment cores would detect and estimate the nature and sources of fluid flow through high permeability pathways in the overriding plate and along the subduction thrust;

characterize the hydrocarbon and gas hydrate system to assist with estimates of heat flow from Bottom Simulating Reflectors (BSR)s, their role in slope stability, and fluid source; and elucidate the response of microbes involved in carbon cycling to changes in methane flux.

The low-energy seismic survey would be collected in a total of 9 grids of intersecting lines of two sizes (see Figure 1 of the IHA application) at exact locations to be determined in the field. The water depths would be very similar to those at the nominal survey locations shown in Figure 1 of the IHA application. The northern and middle sites off the North Island would be the primary study areas, and the southern site off the South Island would be a contingency area that would only be surveyed if time permits. SIO's calculations assume that 7 grids at the primary areas and two grids at the southern site would be surveyed. The total trackline distance of the low-energy seismic survey would be approximately 1,250 km (including the two South Island contingency sites), almost all in water depths greater than 1,000 m.

The procedures to be used for the survey would be similar to those used during previous low-energy seismic surveys by SIO and NSF and would use conventional seismic methodology. The proposed survey would involve one source vessel, the *Revelle*. SIO would deploy a two Sercel Generator Injector (GI) airgun array (each with a discharge volume of 45 in³ [290.3 cm³], in one string, with a total volume of 90 in³ [580.6 cm³]) as an energy source, at a tow depth of up to 2 m (6.6 ft) below the surface (more information on the airguns can be found in SIO's IHA application). The airguns in the array would be spaced approximately 8 m (26.2 ft) apart and 21 m (68.9 ft) astern

of the vessel. The receiving system would consist of one 600 m (1,968.5 ft) long, 48-channel hydrophone streamer(s) towed behind the vessel. Data acquisition is planned along a series of predetermined lines, almost all (approximately 95%) of which would be in water depths greater than 1,000 m. As the GI airguns are towed along the survey lines, the hydrophone streamer would receive the returning acoustic signals and transfer the data to the onboard processing system. The seismic surveys would be conducted while the heat-flow probe is being recharged. All planned seismic data acquisition activities would be conducted by technicians provided by SIO, with onboard assistance by the scientists who have proposed the study. The vessel would be self-contained, and the crew would live aboard the vessel for the entire cruise.

The planned seismic survey (including equipment testing, start-up, line changes, repeat coverage of any areas, and equipment recovery) would consist of approximately 1,250 kilometers (km) (674.9 nautical miles [nmi]) of transect lines (including turns) in the study area in the Southwest Pacific Ocean (see Figures 1 of the IHA application). Approximately 95% of the low-energy seismic survey would occur in water depths greater than 1,000 m. In addition to the operation of the airgun array and heat-flow measurements, a multi-beam echosounder and a sub-bottom profiler would also likely be operated from the *Revelle* continuously throughout the cruise. There would be additional airgun operations associated with equipment testing, ramp-up, and possible line changes or repeat coverage of any areas where initial data quality is sub-standard. In SIO's estimated take calculations, 25% has been added for those additional operations.

TABLE 1—PROPOSED LOW-ENERGY SEISMIC SURVEY ACTIVITIES IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND

Survey length (km)	Total duration (hr) ¹	Airgun array total volume	Time between airgun shots (distance)	Streamer length (m)
1,250 (674.9 nmi)	~135	2 × 45 = 90 in ³ (2 × 1474.8 cm ³)	6 to 10 seconds (18.5 to 31 m or 60.7 to 101.7 ft).	600 (1,968.5 ft)

¹ Airgun operations are planned for no more than approximately 72 continuous hours at a time.

Vessel Specifications

The *Revelle*, a research vessel owned by the U.S. Navy and operated by SIO of the University of California San Diego, would tow the two GI airgun array, as well as the hydrophone streamer. When the *Revelle* is towing

the airgun array and the relatively short hydrophone streamer, the turning rate of the vessel while the gear is deployed is approximately 20 degrees per minute, which is much higher than the limit of 5 degrees per minute for a seismic vessel towing a streamer of more typical length (much greater than 1 km [0.5

nmi]). Thus, the maneuverability of the vessel would not be limited much during operations with the streamer.

The U.S.-flagged vessel, built in 1996, has a length of 83 m (272.3 ft); a beam of 16.0 m (52.5 ft); a maximum draft of 5.2 m (19.5 ft); and a gross tonnage of 3,180. The ship is powered by two 3,000

horsepower (hp) Propulsion General Electric motors) and a 1,180 hp azimuthing jet bowthruster. The GI airgun compressor onboard the vessel is manufactured by Price Air Compressors. The *Revelle's* operation speed during seismic acquisition is typically approximately 9.3 km/hr (5 kts) (varying between 7.4 to 11.1 km/hr [4 to 6 kts]). When not towing seismic survey gear, the *Revelle* typically cruises at 22.2 to 23.1 km/hr (12 to 12.5 kts) and has a maximum speed of 27.8 km/hr (15 kts). The *Revelle* has an operating range of approximately 27,780 km (15,000 nmi) (the distance the vessel can travel without refueling), which is approximately 70 to 75 days. The vessel can accommodate 37 scientists and 22 crew members.

The vessel also has two observation station locations from which Protected Species Observers (PSO) would watch for marine mammals before and during the proposed airgun operations on the *Revelle*. Observing stations would be at the 02 level, with a PSO's eye level approximately 10.4 m (34 ft) above sea level—one forward on the 02 deck commanding a forward-centered, approximately 240° view around the vessel, and one atop the aft hangar, with an aft-centered view that includes the radii around the airguns. The eyes on the bridge watch would be at a height of approximately 15 m (49 ft); PSOs would work on the enclosed bridge and adjoining aft steering station during any inclement weather. More details of the *Revelle* can be found in the IHA application and online at: <https://scripps.ucsd.edu/ships/revelle>.

Acoustic Source Specifications—Seismic Airguns

The *Revelle* would deploy an airgun array, consisting of two 45 in³ Sercel GI airguns as the primary energy source and a 600 m streamer(s) containing hydrophones. The airgun array would have a supply firing pressure of 1,750 pounds per square inch (psi). Seismic pulses for the GI airguns would be emitted at intervals of approximately 6 to 10 seconds. There would be a maximum of approximately 360 shots per hour. The number of shots per hour would vary based upon the vessel speed over ground during the low-energy seismic survey. During firing, a brief (approximately 20 millisecond) pulse sound would be emitted; the airguns would be silent during the intervening periods. The dominant frequency components would range from 0 to 188 Hertz (Hz).

The GI airguns would fire the compressed air volume in unison in “true GI” mode. The GI airguns would

be used in “true GI” mode, that is, the volume of the injector chamber (I) (105 in³ [1721 cm³]) of each GI airgun is greater to that of its generator chamber (G) (45 in³ [737 cm³]) for each airgun. The generator chamber of each GI airgun (45 in³) would be the primary source and the one responsible for introducing the sound pulse into the ocean. The larger (105 in³) injector chamber injects air into the previously-generated bubble to maintain its shape, and would not introduce more sound into the water. The two GI airguns would be spaced approximately 8 m (26.2 ft) apart, side-by-side, 21 m (68.9 ft) behind the *Revelle*, at a depth of up to 2 m during the low-energy seismic survey.

The Nucleus modeling software used at Lamont-Doherty Earth Observatory of Columbia University (L-DEO) does not include GI airguns as part of its airgun library, however signatures and mitigation models have been obtained for two 45 in³ G airguns that are close approximations. For the two 45 in³ airgun array, the source output (downward) is 230.6 dB re 1 μPam 0-to-peak and 235.8 dB re 1 μPam for peak-to-peak. The dominant frequency range would be 0 to 188 Hz for a pair of GI airguns towed at 2 m depth.

During the low-energy seismic survey, the vessel would attempt to maintain a constant cruise speed of approximately 5 knots. The airguns would operate continuously for no more than approximately 72 hours at a time based on operational constraints. The total duration of the airgun operations would not exceed 135 hours. The relatively short, 48-channel hydrophone streamer would provide operational flexibility to allow the low-energy seismic survey to proceed along the designated cruise tracklines. The design of the seismic equipment is to achieve high-resolution images with the ability to correlate to the ultra-high frequency sub-bottom profiling data and provide cross-sectional views to pair with the seafloor bathymetry.

Metrics Used in This Document

This section includes a brief explanation of the sound measurements frequently used in the discussions of acoustic effects in this document. Sound pressure is the sound force per unit area, and is usually measured in micropascals (μPa), where 1 pascal (Pa) is the pressure resulting from a force of one newton exerted over an area of one square meter. Sound pressure level (SPL) is expressed as the ratio of a measured sound pressure and a reference level. The commonly used reference pressure level in underwater acoustics is 1 μPa, and the units for

SPLs are dB re 1 μPa. SPL (in decibels [dB]) = 20 log (pressure/reference pressure).

SPL is an instantaneous measurement and can be expressed as the peak, the peak-to-peak (p-p), or the root mean square (rms). Root mean square, which is the square root of the arithmetic average of the squared instantaneous pressure values, is typically used in discussions of the effects of sounds on vertebrates and all references to SPL in this document refer to the root mean square unless otherwise noted. SPL does not take the duration of a sound into account.

Characteristics of the Airgun Pulses

Airguns function by venting high-pressure air into the water, which creates an air bubble. The pressure signature of an individual airgun consists of a sharp rise and then fall in pressure, followed by several positive and negative pressure excursions caused by the oscillation of the resulting air bubble. The oscillation of the air bubble transmits sounds downward through the seafloor, and the amount of sound transmitted in the near horizontal directions is reduced. However, the airgun array also emits sounds that travel horizontally toward non-target areas.

The nominal downward-directed source levels of the airgun arrays used by SIO on the *Revelle* do not represent actual sound levels that can be measured at any location in the water. Rather, they represent the level that would be found 1 m (3.3 ft) from a hypothetical point source emitting the same total amount of sound as is emitted by the combined GI airguns. The actual received level at any location in the water near the GI airguns would not exceed the source level of the strongest individual source. In this case, that would be about 224.6 dB re 1 μPam peak or 229.8 dB re 1 μPam peak-to-peak for the two 45 in³ airgun array. However, the difference between rms and peak or peak-to-peak values for a given pulse depends on the frequency content and duration of the pulse, among other factors. Actual levels experienced by any organism more than 1 m from either GI airgun would be significantly lower.

Accordingly, L-DEO has predicted and modeled the received sound levels in relation to distance and direction from the two GI airgun array. These are the nominal source levels applicable to downward propagation. A detailed description of L-DEO's modeling for this survey's marine seismic source arrays for protected species mitigation is provided in the “Programmatic

Environmental Impact Statement/ Overseas Environmental Impact Statement prepared for Marine Seismic Research that is funded by the National Science Foundation and conducted by the U.S. Geological Survey” (NSF/USGS PEIS, 2011). The NSF/USGS PEIS discusses the characteristics of the airgun pulses. NMFS refers the reviewers to that document for additional information.

Predicted Sound Levels for the Airguns

To estimate takes and determine mitigation (i.e., buffer and exclusion) zones for the airgun array to be used, received sound levels have been modeled by L-DEO for a number of airgun configurations, including two 45 in³ G airguns, in relation to distance and direction from the airguns (see Figure 2 of the IHA application). The model does not allow for bottom interactions, and is most directly applicable to deep water. Because the model results are for G airguns, which have more energy than GI airguns of the same size, those distances overestimate (by approximately 10%) the distances for the two 45 in³ GI airguns. Although the distances are overestimated, no adjustments for this have been made to the radii distances in Table 2 (below). Based on the modeling, estimates of the maximum distances from the GI airguns where sound levels of 190, 180, and 160 dB re 1 μPa (rms) are predicted to be received in intermediate and deep water

are shown in Table 2 (see Table 1 of the IHA application).

Empirical data concerning the 190, 180, and 160 dB (rms) distances were acquired for various airgun arrays based on measurements during the acoustic verification studies conducted by L-DEO in the northern Gulf of Mexico (GOM) in 2003 (Tolstoy *et al.*, 2004) and 2007 to 2008 (Tolstoy *et al.*, 2009; Diebold *et al.*, 2010). Results of the 18 and 36 airgun array are not relevant for the two GI airguns to be used in the proposed low-energy seismic survey because the airgun arrays are not the same size or volume. The empirical data for the 6, 10, 12, and 20 airgun arrays indicate that, for deep water, the L-DEO model tends to overestimate the received sound levels at a given distance (Tolstoy *et al.*, 2004). For the two G airgun array, measurements were obtained only in shallow water. When compared to measurements in acquired in deep water, mitigation radii provided by the L-DEO model for the proposed airgun operations were found to be conservative. The acoustic verification surveys also showed that distances to given received levels vary with water depth; these are larger in shallow water, while intermediate/slope environments show characteristics intermediate between those of shallow water and those of deep water environments, and documented the influence of a sloping seafloor. The only measurements obtained for intermediate depths during either survey were for the 36-airgun

array in 2007 to 2008 (Diebold *et al.*, 2010). Following results obtained at this site and earlier practice, a correction factor of 1.5, irrespective of distance to the airgun array, is used to derive intermediate-water radii from modeled deep-water radii.

Measurements were not made for a two GI airgun array in intermediate and deep water; however, SIO proposes to use the buffer and exclusion zones predicted by L-DEO’s model for the proposed GI airgun operations in intermediate and deep water, although they are likely conservative given the empirical results for the other arrays. Using the L-DEO model, Table 2 (below) shows the distances at which three rms sound levels are expected to be received from the two GI airguns. The 160 dB re 1 μPam (rms) isopleth is the threshold specified by NMFS for potential Level B (behavioral) harassment from impulsive noise for both cetaceans and pinnipeds. The 180 and 190 dB re 1 μPam (rms) isopleths are the thresholds currently used to estimate potential Level A harassment as specified by NMFS (2000) and are applicable to cetaceans and pinnipeds, respectively. Table 2 summarizes the predicted distances at which sound levels (160, 180, and 190 dB [rms]) are expected to be received from the two airgun array (each 45 in³) operating in intermediate water (100 to 1,000 m [328.1 to 3,280 ft]) and deep water (>1,000 m) depths.

TABLE 2—PREDICTED AND MODELED (TWO 45 IN³ GI AIRGUN ARRAY) DISTANCES TO WHICH SOUND LEVELS ≥160, 180, AND 190 dB RE 1 μPA (rms) COULD BE RECEIVED IN INTERMEDIATE AND DEEP WATER DURING THE PROPOSED LOW-ENERGY SEISMIC SURVEY IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND, MAY TO JUNE 2015

Source and total volume	Tow depth (m)	Water depth (m)	Predicted RMS radii distances (m) for 2 GI airgun array		
			160 dB	180 dB	190 dB
Two 45 in ³ GI Airguns (90 in ³).	2	Intermediate (100 to 1,000).	600 (1,968.5 ft)	100 (328.1 ft)	15 (49.2 ft) *100 would be used for pinnipeds as described in NSF/USGS PEIS*
Two 45 in ³ GI Airguns (90 in ³).	2	Deep (>1,000)	400 (1,312.3 ft)	100 (328.1 m)	10 (32.8 ft) *100 would be used for pinnipeds as described in NSF/USGS PEIS*

Based on the NSF/USGS PEIS and Record of Decision, for situations which incidental take of marine mammals is anticipated, proposed exclusion zones of 100 m for cetaceans and pinnipeds for all low-energy acoustic sources in water depths greater than 100 m would be implemented.

NMFS expects that acoustic stimuli resulting from the proposed operation of the two GI airgun array has the potential to harass marine mammals. NMFS does not expect that the movement of the *Revelle*, during the conduct of the low-energy seismic survey, has the potential

to harass marine mammals because the relatively slow operation speed of the vessel (approximately 5 kts; 9.3 km/hr; 5.8 mph) during seismic data acquisition should allow marine mammals to avoid the vessel.

Bathymetric Survey

Along with the low-energy airgun operations, two additional geophysical (detailed swath bathymetry) measurements focused on a specific study area within the Southwest Pacific Ocean would be made using hull-mounted sonar system instruments from

the *Revelle* for operational and navigational purposes. The ocean floor would be mapped with the Kongsberg EM 122 multi-beam echosounder and a Knudsen Chirp 3260 sub-bottom profiler. During bathymetric survey operations, when the vessel is not towing seismic equipment, its average speed would be approximately 10.1 kts (18.8 km/hr). In cases where higher resolution bathymetric data is sought, the average speed may be as low as 5 kts (9.3 km/hr). These sound sources would be operated continuously from the *Revelle* throughout the cruise. Operating

characteristics for the instruments to be used are described below.

Multi-Beam Echosounder (Kongsberg EM 122)—The hull-mounted multi-beam sonar would be operated continuously during the cruise to map the ocean floor. This instrument would operate at a frequency of 10.5 to 13 (usually 12) kilohertz (kHz) and would be hull-mounted. The transmitting beamwidth would be 1 or 2° fore to aft and 150° athwartship (cross-track). The estimated maximum source energy level would be 242 dB re 1 μ Pa (rms). Each ‘ping’ of eight (in water greater than 1,000 m or four (in water less than 1,000 m) successive fan-shaped transmissions, each ensonifying a sector that extends 1° fore to aft. Continuous-wave signals increase from 2 to 15 milliseconds (ms) in water depths up to 2,600 m (8,530 ft), and FM chirp signals up to 100 ms long would be used in water greater than 2,600 m. The successive transmission span an overall cross-track angular extent of about 150°, with 2 ms gaps between the pings for successive sectors.

Sub-Bottom Profiler—The *Revelle* would operate a Knudsen 3260 sub-bottom profiler continuously throughout the cruise simultaneously to map and provide information about the seafloor sedimentary features and bottom topography that is mapped simultaneously with the multi-beam echosounder. The beam of the sub-bottom profiler would be transmitted as a 27° cone, directed downward by a 3.5 kHz transducer in the hull of the *Revelle*. The nominal power output would be 10 kilowatt (kW), but the actual maximum radiated power would be 3 kW or 222 dB (rms). The ping duration would be up to 64 ms, and the ping interval would be 1 second. A common mode of operation is a broadcast five pulses at 1 second intervals followed by a 5 second pause. The sub-bottom profiler would be capable of reaching depths of 10,000 m (32,808.4 ft).

Acoustic Locator (Pinger)—A pinger would be deployed with certain instruments and equipment (e.g., heat-flow probe) so these devices can be located in the event they become detached from their lines. The pinger used in the heat-flow measurement activities would be the Datasonics model BFP-312HP. A pinger typically operates at a frequency of 32.8 kHz, generates a 5 ms pulse per second (10 pulses over a 10 second period), and has an acoustical output of 210 dB re 1 μ Pa (rms). The pinger would be used during heat-flow measurement operations only. It would operate continuously during each heat-flow probe deployment. Each

heat-flow probe measurement would last approximately 24 hours.

Heat-Flow Probe Deployment

Heat-flow measurements would be made using a “violin-bow” probe with 11 thermistors that provides real time (analog) telemetry of the thermal gradient and in-situ thermal conductivity. The heat-flow probe that would be used on the *Revelle* consists of a lance 6 centimeter (cm) (2.4 in) in diameter and 3.5 m (11.5 ft) long, a sensor tube housing thermistors and heater wires, and a 560 kg (1,234.6 lb) weight stand. The probe would be lowered to the bottom, and a 12 kHz pinger attached to the wire approximately 50 m (164 ft) above the instrument would monitor the distance between the probe and bottom. The probe would be driven into the sediment by gravity, and temperatures within the sediment would be measured with equally spaced thermistors. On completion of a measurement, the instrument would be hoisted 100 to 500 m (328.1 to 1,640.4 ft) above the sediment, the ship is maneuvered to a new position, and the process is repeated. Heat-flow measurements can generally be made at a rate of 1 to 2 hours per measurement, approximately 15 minutes for the actual measurement and 45 to 90 minutes to reposition the ship and probe. Internal power allows 20 to 24 measurements during a single lowering of the tool, with profiles lasting as long as 48 hours. Proposed heat-flow measurements would have a nominal spacing of 0.5 to 1 km (0.3 to 0.5 nmi), which would be decreased in areas of significant basement relief or of large changes in gradient. Heat flow transect locations are shown in Figure 1 of the IHA application, and details of the probe and its deployment are given in Section (f) of the IHA application. In total, approximately 200 heat-flow measurements would be made.

Description of the Marine Mammals in the Specified Geographic Area of the Proposed Specified Activity

Few scientific systematic surveys for marine mammals have been conducted in the waters of New Zealand, and these mainly consist of single-species surveys in shallow coastal waters (e.g., Dawson *et al.*, 2004; Slooten *et al.*, 2004, 2006). Large-scale, multi-species marine mammal surveys are lacking. Various sources for data on sightings in the proposed study area were used to describe the occurrence of marine mammals in the waters of New Zealand, such as opportunistic sighting records presented in previous reports (including the New Zealand Department of

Conservation marine mammals sighting database) considered in evaluating potential marine mammals in the proposed action area.

New Zealand is considered a “hotspot” for marine mammal species richness (Kaschner *et al.*, 2011). The marine mammals that generally occur in the proposed action area belong to three taxonomic groups: Mysticetes (baleen whales), odontocetes (toothed whales), and pinnipeds (seals and sea lions). The marine mammal species that could potentially occur within the Southwest Pacific Ocean in proximity to the proposed action area East of New Zealand include 30 species of cetaceans (21 odontocetes and 9 mysticetes) and 2 species of pinnipeds (32 total species of marine mammals).

Marine mammal species likely to be encountered in the proposed study area that are listed as endangered under the U.S. Endangered Species Act of 1973 (ESA; 16 U.S.C. 1531 *et seq.*), includes the southern right (*Eubalaena australis*), humpback (*Megaptera novaeangliae*), sei (*Balaenoptera borealis*), fin (*Balaenoptera physalus*), blue (*Balaenoptera musculus*), and sperm (*Physeter macrocephalus*) whale. The Maui’s dolphin (*Cephalorhynchus hectori maui*) and New Zealand sea lion (*Phocartos hookeri*) are two other species are ranked as “nationally critical” in New Zealand (Baker *et al.*, 2010). Maui’s dolphin is only found along the west coast of the North Island. The northern range of the New Zealand sea lion is not expected to extend to the proposed study area based on New Zealand’s National Aquatic Biodiversity Information System (NABIS, 2014) and is not considered further.

In addition to the marine mammal species known to occur in the Southwest Pacific Ocean off the east coast of New Zealand, there are 18 species of marine mammals (12 cetacean and 6 pinniped species) with ranges that are known to potentially occur in the waters of the proposed study area, but they are categorized as “vagrant” under the New Zealand Threat Classification System (Baker *et al.*, 2010). These include: Dwarf sperm whale (*Kogia sima*), Arnoux’s beaked whale (*Berardius arnouxii*), ginkgo-toothed beaked whale (*Mesoplodon ginkgodens*), pygmy beaked whale (*Mesoplodon peruvianis*), Type B, C, and D killer whale (*Orcinus orca*), melon-headed whale (*Peponocephala electra*), Risso’s dolphin (*Grampus griseus*), Fraser’s dolphin (*Lagenodelphis hosei*), pantropical spotted dolphin (*Stenella attenuata*), striped dolphin (*Stenella coeruleoalba*), rough-toothed dolphin (*Steno bredanensis*), spectacled

porpoise (*Phocoena dioptrica*), Antarctic fur seal (*Arctocephalus gazelle*), Subantarctic fur seal (*Arctocephalus tropicalis*), crabeater seal (*Lobodon carcinophagus*), leopard seal (*Hydrurga leptonyx*), Ross seal (*Ommatophoca rossi*), and Weddell seal (*Leptonychotes weddellii*). According to Jefferson *et al.* (2008), the distributional range of Hubb's beaked whale (*Mesoplodon carlhubbsi*) and True's

beaked whale (*Mesoplodon mirus*) may also include New Zealand waters. There are no records of Hubb's beaked whale in New Zealand, and only a single record of True's beaked whale, which stranded on the west coast of South Island in November 2011 (Constantine *et al.*, 2014). The spinner dolphin's (*Stenella longirostris*) range includes tropical and subtropical zones 40° North to 40° South, but would be considered

vagrant as well. However, these species are not expected to occur where the proposed activities would take place. These species are not considered further in this document. Table 3 (below) presents information on the habitat, occurrence, distribution, abundance, population, and conservation status of the species of marine mammals that may occur in the proposed study area during May to June 2015.

TABLE 3—THE HABITAT, OCCURRENCE, RANGE, REGIONAL ABUNDANCE, AND CONSERVATION STATUS OF MARINE MAMMALS THAT MAY OCCUR IN OR NEAR THE PROPOSED LOW-ENERGY SEISMIC SURVEY AREA IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND

[See text and tables 2 in SIO's IHA application for further details]

Species	Habitat	Occurrence	Range	Population estimate	ESA ¹	MMPA ²
Mysticetes						
Southern right whale (<i>Eubalaena australis</i>).	Coastal, shelf, pelagic.	Common	Circumpolar 20 to 55° South.	8,000 ³ to 15,000 ⁴ —Worldwide 12,000 ¹² —Southern Hemisphere 2,700 ¹² —Sub-Antarctic New Zealand.	EN	D
Pygmy right whale (<i>Caperea marginata</i>).	Pelagic and coastal.	Rare	Circumpolar 30 to 55° South.	NA	NL	NC
Humpback whale (<i>Megaptera novaeangliae</i>).	Pelagic, nearshore waters, and banks.	Common	Cosmopolitan Migratory ...	35,000 to 42,000 ^{3 12} —Southern Hemisphere.	EN	D
Minke whale (<i>Balaenoptera acutorostrata</i> including dwarf sub-species).	Pelagic and coastal.	Uncommon ..	Circumpolar—Southern Hemisphere to 65° South.	720,000 to 750,000 ^{12 14 15} —Southern Hemisphere.	NL	NC
Antarctic minke whale (<i>Balaenoptera bonaerensis</i>).	Pelagic, ice floes, coastal.	Uncommon ..	7° South to ice edge (usually 20 to 65° South).	720,000 to 750,000 ^{12 14 15} —Southern Hemisphere.	NL	NC
Bryde's whale (<i>Balaenoptera edeni</i>).	Pelagic and coastal.	Rare	Circumglobal—Tropical and Subtropical Zones.	At least 30,000 to 40,000 ³ —Worldwide 21,000 ¹² —Northwestern Pacific Ocean 48,109 ¹³ .	NL	NC
Sei whale (<i>Balaenoptera borealis</i>).	Primarily offshore, pelagic.	Uncommon ..	Migratory, Feeding Concentration 40 to 50° South.	80,000 ³ —Worldwide 10,000 ¹⁴ —South of Antarctic Convergence.	EN	D
Fin whale (<i>Balaenoptera physalus</i>).	Continental slope, pelagic.	Uncommon ..	Cosmopolitan, Migratory ..	140,000 ³ —Worldwide 15,000 ¹⁴ —South of Antarctic Convergence.	EN	D
Blue whale (<i>Balaenoptera musculus</i> ; including pygmy blue whale [<i>Balaenoptera musculus breviceauda</i>]).	Pelagic, shelf, coastal.	Uncommon ..	Migratory Pygmy blue whale—North of Antarctic Convergence 55° South.	8,000 to 9,000 ³ —Worldwide 2,300 ¹² —True Southern Hemisphere 1,500 ¹⁴ —Pygmy.	EN	D
Odontocetes						
Sperm whale (<i>Physeter macrocephalus</i>).	Pelagic, deep sea	Common	Cosmopolitan, Migratory ...	360,000 ³ —Worldwide 30,000 ¹³ —South of Antarctic Convergence.	EN	D
Dwarf sperm whale (<i>Kogia sima</i>).	Shelf, Pelagic	Vagrant	Circumglobal—Tropical and Temperate Zones.	NA	NL	NC
Pygmy sperm whale (<i>Kogia breviceps</i>).	Shelf, Pelagic	Uncommon ..	Circumglobal—Temperate Zones.	NA	NL	NC
Arnoux's beaked whale (<i>Berardius arnuxii</i>).	Pelagic	Vagrant	Circumpolar in Southern Hemisphere, 24 to 78° South.	NA	NL	NC
Cuvier's beaked whale (<i>Ziphius cavirostris</i>).	Pelagic	Uncommon ..	Cosmopolitan	600,000 ^{14 16}	NL	NC
Southern bottlenose whale (<i>Hyperoodon planifrons</i>).	Pelagic	Rare	Circumpolar—30° South to ice edge.	500,000 ³ —South of Antarctic Convergence 600,000 ^{14 16} .	NL	NC
Shepherd's beaked whale (<i>Tasmacetus shepherdi</i>).	Pelagic	Rare	Circumpolar—Cold temperate waters Southern Hemisphere.	600,000 ^{14 16}	NL	NC

TABLE 3—THE HABITAT, OCCURRENCE, RANGE, REGIONAL ABUNDANCE, AND CONSERVATION STATUS OF MARINE MAMMALS THAT MAY OCCUR IN OR NEAR THE PROPOSED LOW-ENERGY SEISMIC SURVEY AREA IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND—Continued

[See text and tables 2 in SIO's IHA application for further details]

Species	Habitat	Occurrence	Range	Population estimate	ESA ¹	MMPA ²
Andrew's beaked whale (<i>Mesoplodon bowdoini</i>).	Pelagic	Rare	Circumpolar—temperate waters of Southern Hemisphere, 32 to 55° South.	600,000 ^{14 16}	NL	NC
Blainville's beaked whale (<i>Mesoplodon densirostris</i>).	Pelagic	Rare	Circumglobal—tropical and temperate waters.	600,000 ^{14 16}	NL	NC
Ginkgo-toothed beaked whale (<i>Mesoplodon ginkgodens</i>).	Pelagic	Vagrant	Tropical and Temperate waters—Indo-Pacific Ocean.	NA	NL	NC
Gray's beaked whale (<i>Mesoplodon grayi</i>).	Pelagic	Common	30° South to Antarctic waters.	600,000 ^{14 16}	NL	NC
Hector's beaked whale (<i>Mesoplodon hectori</i>).	Pelagic	Rare	Circumpolar—cool temperate waters of Southern Hemisphere.	600,000 ^{14 16}	NL	NC
Hubb's beaked whale (<i>Mesoplodon carlhubbsi</i>).	Pelagic	Vagrant	North Pacific Ocean	NA	NL	NC
Pygmy beaked whale (<i>Mesoplodon peruvianis</i>).	Pelagic	Vagrant	28° North to 30° South in Pacific Ocean.	NA	NL	NC
Spade-toothed beaked whale (<i>Mesoplodon traversii</i>).	Pelagic	Rare	Circumantarctic	600,000 ^{14 16}	NL	NC
Strap-toothed beaked whale (<i>Mesoplodon layardii</i>).	Pelagic	Uncommon ..	30° South to Antarctic Convergence.	600,000 ^{14 16}	NL	NC
True's beaked whale (<i>Mesoplodon mirus</i>).	Pelagic	Vagrant	Anti-tropical in Northern and Southern Hemisphere.	NA	NL	NC
Killer whale (<i>Orcinus orca</i>)	Pelagic, shelf, coastal, pack ice.	Common	Cosmopolitan	80,000 ³ —South of Antarctic Convergence.	NL	NC
False killer whale (<i>Pseudorca crassidens</i>).	Pelagic, shelf, coastal.	Uncommon ..	Circumglobal—tropical and warmer temperate water.	NA	NL	NC
Long-finned pilot whale (<i>Globicephala melas</i>).	Pelagic, shelf, coastal.	Common	Circumpolar—19 to 68° South in Southern Hemisphere.	200,000 ^{3 5 14} —South of Antarctic Convergence.	NL	NC
Short-finned pilot whale (<i>Globicephala macrocephalus</i>).	Pelagic, shelf, coastal.	Uncommon ..	Circumglobal—50° North to 40° South.	At least 600,000 ³ —Worldwide.	NL	NC
Melon-headed whale (<i>Peponocephala electra</i>).	Pelagic, shelf, coastal.	Vagrant	Circumglobal—40° North to 35° South.	45,000 ³ —Eastern Tropical Pacific Ocean.	NL	NC
Bottlenose dolphin (<i>Tursiops truncatus</i>).	Coastal, shelf, off-shore.	Common	45° North to 45° South	At least 614,000 ³ —Worldwide.	NL, *C	NC
Dusky dolphin (<i>Lagenorhynchus obscurus</i>).	Shelf, slope	Common	Temperate waters—Southern Hemisphere.	12,000 to 20,000 ¹⁷ —New Zealand.	NL	NC
Fraser's dolphin (<i>Lagenodelphis hosei</i>).	Pelagic	Vagrant	Pantropical—30° North to 30° South.	289,000 ³ —Eastern Tropical Pacific Ocean.	NL	NC
Hector's dolphin (<i>Cephalorhynchus hectori</i> ; including Maui's dolphin subspecies [<i>C. h. mauii</i>]).	Nearshore	Rare	Shallow coastal waters—New Zealand (Maui's dolphin—west North Island).	7,400 ¹⁷	C	NC
Hourglass dolphin (<i>Lagenorhynchus cruciger</i>).	Pelagic, ice edge	Uncommon ..	33° South to pack ice	144,000 ³ to 150,000 ¹⁴ —South of Antarctic Convergence.	NL	NC
Pantropical spotted dolphin (<i>Stenella attenuata</i>).	Coastal, shelf, slope.	Vagrant	Circumglobal—40° North to 40° South.	At least 2,000,000 ³ —Worldwide.	NL	NC
Spinner dolphin (<i>Stenella longirostris</i>).	Mainly nearshore	Vagrant	Circumglobal—40° North to 40° South.	At least 1,200,000 ³ —Worldwide.	NL	NC
Striped dolphin (<i>Stenella coeruleoalba</i>).	Off continental shelf, convergence zones, upwelling.	Vagrant	Circumglobal—50 to 40 South.	At least 1,100,000 ³ —Worldwide.	NL	NC
Risso's dolphin (<i>Grampus griseus</i>).	Slope, Pelagic	Vagrant	Circumglobal—Tropical and Temperate waters.	At least 330,000 ³ —Worldwide.	NL	NC
Rough-toothed dolphin (<i>Steno bredanensis</i>).	Pelagic	Vagrant	Circumglobal—40° North to 35° South.	NA	NL	NC

TABLE 3—THE HABITAT, OCCURRENCE, RANGE, REGIONAL ABUNDANCE, AND CONSERVATION STATUS OF MARINE MAMMALS THAT MAY OCCUR IN OR NEAR THE PROPOSED LOW-ENERGY SEISMIC SURVEY AREA IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND—Continued

[See text and tables 2 in SIO's IHA application for further details]

Species	Habitat	Occurrence	Range	Population estimate	ESA ¹	MMPA ²
Short-beaked common dolphin (<i>Delphinus delphis</i>).	Pelagic	Common	Circumglobal—tropical and warm temperate waters.	At least 3,500,000 ³ —Worldwide.	NL	NC
Southern right whale dolphin (<i>Lissodelphis peronii</i>).	Pelagic	Uncommon ..	12 to 65° South	NA	NL	NC
Spectacled porpoise (<i>Phocoena dioptrica</i>).	Coastal, pelagic ...	Vagrant	Circumpolar—Southern Hemisphere.	NA	NL	NC
Pinnipeds						
Crabeater seal (<i>Lobodon carcinophaga</i>).	Coastal, pack ice	Vagrant	Circumpolar—Antarctic	5,000,000 to 15,000,000 ^{3,6} —Worldwide.	NL	NC
Leopard seal (<i>Hydrurga leptonyx</i>).	Pack ice, sub-Antarctic islands.	Vagrant	Sub-Antarctic islands to pack ice.	220,000 to 440,000 ^{3,7} —Worldwide.	NL	NC
Ross seal (<i>Ommatophoca rossii</i>).	Pack ice, smooth ice floes, pelagic.	Vagrant	Circumpolar—Antarctic	130,000 ³ 20,000 to 220,000 ¹¹ —Worldwide.	NL	NC
Weddell seal (<i>Leptonychotes weddellii</i>).	Fast ice, pack ice, sub-Antarctic islands.	Vagrant	Circumpolar—Southern Hemisphere.	500,000 to 1,000,000 ^{3,8} —Worldwide.	NL	NC
Southern elephant seal (<i>Mirounga leonina</i>).	Coastal, pelagic, sub-Antarctic waters.	Uncommon ..	Circumpolar—Antarctic Convergence to pack ice.	640,000 ⁹ to 650,000 ³ —Worldwide 470,000—South Georgia Island ¹¹ 607,000 ¹⁷ .	NL	NC
Antarctic fur seal (<i>Arctocephalus gazella</i>).	Shelf, rocky habitats.	Vagrant	Sub-Antarctic islands to pack ice edge.	1,600,000 ¹⁰ to 3,000,000 ³ —Worldwide.	NL	NC
New Zealand fur seal (<i>Arctocephalus forsteri</i>).	Rocky habitats, sub-Antarctic islands.	Common	North and South Islands, New Zealand Southern and Western Australia.	135,000 ³ —Worldwide 50,000 to 100,000 ¹⁸ —New Zealand.	NL	NC
Subantarctic fur seal (<i>Arctocephalus tropicalis</i>).	Shelf, rocky habitats.	Vagrant	Subtropical front to sub-Antarctic islands and Antarctica.	Greater than 310,000 ³ —Worldwide.	NL	NC
New Zealand sea lion (<i>Phocarctos hookeri</i>).	Shelf, rocky habitats.	Rare	Sub-Antarctic islands south of New Zealand.	12,500 ³	NL	NC

NA = Not available or not assessed.

³ Fjordland population.

¹ U.S. Endangered Species Act: EN = Endangered, T = Threatened, DL = Delisted, NL = Not listed, C = Candidate.

² U.S. Marine Mammal Protection Act: D = Depleted, S = Strategic, NC = Not Classified.

³ Jefferson *et al.*, 2008.

⁴ Kenney, 2009.

⁵ Olson, 2009.

⁶ Bengston, 2009.

⁷ Rogers, 2009.

⁸ Thomas and Terhune, 2009.

⁹ Hindell and Perrin, 2009.

¹⁰ Arnould, 2009.

¹¹ Academic Press, 2009.

¹² IWC, 2014.

¹³ IWC, 1981.

¹⁴ Boyd, 2002.

¹⁵ Dwarf and Antarctic minke whale combined.

¹⁶ All Antarctic beaked whales combined.

¹⁷ New Zealand Department of Conservation.

¹⁸ Suisted and Neale, 2004.

Refer to sections 3 and 4 of SIO's IHA application for detailed information regarding the abundance and distribution, population status, and life history and behavior of these marine mammal species and their occurrence in the proposed action area. The IHA application also presents how SIO calculated the estimated densities for the marine mammals in the proposed study area. NMFS has reviewed these

data and determined them to be the best available scientific information for the purposes of the proposed IHA.

Potential Effects of the Proposed Specified Activity on Marine Mammals

This section includes a summary and discussion of the ways that the types of stressors associated with the specified activity (*e.g.*, seismic airgun operation, vessel movement, and gear deployment)

have been thought to impact marine mammals. This discussion may also include reactions that we consider to rise to the level of a take and those that we do not consider to rise to the level of take (for example, with acoustics, we may include a discussion of studies that showed animals not reacting at all to sound or exhibiting barely measurable avoidance). This section is intended as a background of potential effects and

does not consider either the specific manner in which this activity would be carried out or the mitigation that would be implemented, and how either of those would shape the anticipated impacts from this specific activity. The “Estimated Take by Incidental Harassment” section later in this document would include a quantitative analysis of the number of individuals that are expected to be taken by this activity. The “Negligible Impact Analysis” section will include the analysis of how this specific activity will impact marine mammals and will consider the content of this section, the “Estimated Take by Incidental Harassment” section, the “Proposed Mitigation” section, and the “Anticipated Effects on Marine Mammal Habitat” section to draw conclusions regarding the likely impacts of this activity on the reproductive success or survivorship of individuals and from that on the affected marine mammal populations or stocks.

When considering the influence of various kinds of sound on the marine environment, it is necessary to understand that different kinds of marine life are sensitive to different frequencies of sound. Based on available behavioral data, audiograms have been derived using auditory evoked potentials, anatomical modeling, and other data, Southall *et al.* (2007) designate “functional hearing groups” for marine mammals and estimate the lower and upper frequencies of functional hearing of the groups. The functional groups and the associated frequencies are indicated below (though animals are less sensitive to sounds at the outer edge of their functional range and most sensitive to sounds of frequencies within a smaller range somewhere in the middle of their functional hearing range):

- Low-frequency cetaceans (13 species of mysticetes): Functional hearing is estimated to occur between approximately 7 Hz and 30 kHz;

- Mid-frequency cetaceans (32 species of dolphins, six species of larger toothed whales, and 19 species of beaked and bottlenose whales):

Functional hearing is estimated to occur between approximately 150 Hz and 160 kHz;

- High-frequency cetaceans (eight species of true porpoises, six species of river dolphins, *Kogia* spp., the franciscana [*Pontoporia blainvillei*], and four species of cephalorhynchids):

Functional hearing is estimated to occur between approximately 200 Hz and 180 kHz; and

- Phocid pinnipeds in water: Functional hearing is estimated to occur

between approximately 75 Hz and 100 kHz;

- Otariid pinnipeds in water: Functional hearing is estimated to occur between approximately 100 Hz and 40 kHz.

As mentioned previously in this document, 32 marine mammal species (30 cetacean and 2 pinniped species) are likely to occur in the proposed low-energy seismic survey area. Of the 30 cetacean species likely to occur in SIO’s proposed action area, 9 are classified as low-frequency cetaceans (southern right, pygmy right, humpback, minke, Antarctic minke, Bryde’s, sei, fin, and blue whale), 20 are classified as mid-frequency cetaceans (sperm, Cuvier’s beaked, Shepherd’s beaked, southern bottlenose, Andrew’s beaked, Blainville’s beaked, Gray’s beaked, Hector’s beaked, spade-toothed beaked, strap-toothed beaked, killer, false killer, long-finned pilot, and short-finned pilot whale, and bottlenose, dusky, Hector’s, hourglass, short-beaked common, and southern right whale dolphin), and 1 is classified as high-frequency cetaceans (pygmy sperm whale) (Southall *et al.*, 2007). Of the 2 pinniped species likely to occur in SIO’s proposed action area, 1 is classified as phocid (southern elephant seal) and 1 is classified as otariid (New Zealand fur seal) (Southall *et al.*, 2007). A species functional hearing group is a consideration when we analyze the effects of exposure to sound on marine mammals.

Acoustic stimuli generated by the operation of the airguns, which introduce sound into the marine environment, have the potential to cause Level B harassment of marine mammals in the proposed study area. The effects of sounds from airgun operations might include one or more of the following: Tolerance, masking of natural sounds, behavioral disturbance, temporary or permanent hearing impairment, or non-auditory physical or physiological effects (Richardson *et al.*, 1995; Gordon *et al.*, 2004; Nowacek *et al.*, 2007; Southall *et al.*, 2007). Permanent hearing impairment, in the unlikely event that it occurred, would constitute injury, but temporary threshold shift (TTS) is not an injury (Southall *et al.*, 2007; Le Prell, 2012). Although the possibility cannot be entirely excluded, it is unlikely that the proposed project would result in any cases of temporary or permanent hearing impairment, or any significant non-auditory physical or physiological effects. Based on the available data and studies described here, some behavioral disturbance is expected. A more comprehensive review of these issues can be found in the NSF/USGS PEIS

(2011) and L-DEO’s “Final Environmental Assessment of a Marine Geophysical Survey by the R/V *Marcus G. Langseth* in the Atlantic Ocean off Cape Hatteras, September to October 2014.”

Tolerance

Richardson *et al.* (1995) defines tolerance as the occurrence of marine mammals in areas where they are exposed to human activities or man-made noise. In many cases, tolerance develops by the animal habituating to the stimulus (*i.e.*, the gradual waning of responses to a repeated or ongoing stimulus) (Richardson *et al.*, 1995; Thorpe, 1963), but because of ecological or physiological requirements, many marine animals may need to remain in areas where they are exposed to chronic stimuli (Richardson *et al.*, 1995).

Numerous studies have shown that pulsed sounds from airguns are often readily detectable in the water at distances of many kilometers (Nieukirk *et al.*, 2012). Several studies have shown that marine mammals at distances more than a few kilometers from operating seismic vessels often show no apparent response. That is often true even in cases when the pulsed sounds must be readily audible to the animals based on measured received levels and the hearing sensitivity of the marine mammal group. Although various baleen whales and toothed whales, and (less frequently) pinnipeds have been shown to react behaviorally to airgun pulses under some conditions, at other times marine mammals of all three types have shown no overt reactions. The relative responsiveness of baleen and toothed whales are quite variable.

Masking

The term masking refers to the inability of a subject to recognize the occurrence of an acoustic stimulus as a result of the interference of another acoustic stimulus (Clark *et al.*, 2009). Introduced underwater sound may, through masking, reduce the effective communication distance of a marine mammal species if the frequency of the source is close to that used as a signal by the marine mammal, and if the anthropogenic sound is present for a significant fraction of the time (Richardson *et al.*, 1995).

The airguns for the proposed low-energy seismic survey have dominant frequency components of 0 to 188 Hz. This frequency range fully overlaps the lower part of the frequency range of odontocete calls and/or functional hearing (full range about 150 Hz to 180 kHz). Airguns also produce a small portion of their sound at mid and high

frequencies that overlap most, if not all, frequencies produced by odontocetes. While it is assumed that mysticetes can detect acoustic impulses from airguns and vessel sounds (Richardson *et al.*, 1995a), sub-bottom profilers, and most of the multi-beam echosounders would likely be detectable by some mysticetes based on presumed mysticete hearing sensitivity. Odontocetes are presumably more sensitive to mid to high frequencies produced by the multi-beam echosounders and sub-bottom profilers than to the dominant low frequencies produced by the airguns and vessel. A more comprehensive review of the relevant background information for odontocetes appears in Section 3.6.4.3, Section 3.7.4.3 and Appendix E of the NSF/USGS PEIS (2011).

Masking effects of pulsed sounds (even from large arrays of airguns) on marine mammal calls and other natural sounds are expected to be limited. Because of the intermittent nature and low duty cycle of seismic airgun pulses, animals can emit and receive sounds in the relatively quiet intervals between pulses. However, in some situations, reverberation occurs for much or the entire interval between pulses (*e.g.*, Simard *et al.*, 2005; Clark and Gagnon, 2006) which could mask calls. Situations with prolonged strong reverberation are infrequent. However, it is common for reverberation to cause some lesser degree of elevation of the background level between airgun pulses (Gedamke, 2011; Guerra *et al.*, 2011, 2013), and this weaker reverberation presumably reduces the detection range of calls and other natural sound to some degree. Guerra *et al.* (2013) reported that ambient noise levels between seismic pulses were elevated because of reverberation at ranges of 50 km (27 nmi) from the seismic source. Based on measurements in deep water of the Southern Ocean, Gedamke (2011) estimated that the slight elevation of background levels during intervals between pulses reduced blue and fin whale communication space by as much as 36 to 51% when a seismic survey was operating 450 to 2,800 km (243 to 1,511.9 nmi) away. Based on preliminary modeling, Wittekind *et al.* (2013) reported that airgun sounds could reduce the communication range of blue and fin whales 2,000 km (1,079.9 nmi) from the seismic source. Klinck *et al.* (2012) also found reverberation effects between pulses. Nieu Kirk *et al.* (2012) and Blackwell *et al.* (2013) noted the potential for masking effects from seismic surveys on large whales.

Some baleen and toothed whales are known to continue calling in the presence of seismic pulses, and their

calls can usually be heard between the seismic pulses (*e.g.*, Richardson *et al.*, 1986; McDonald *et al.*, 1995; Greene *et al.*, 1999; Nieu Kirk *et al.*, 2004, 2012; Smultea *et al.*, 2004; Holst *et al.*, 2005a,b, 2006; and Dunn and Hernandez, 2009). However, Clark and Gagnon (2006) reported that fin whales in the North Atlantic Ocean went silent for an extended period starting soon after the onset of a seismic survey in the area. Similarly, there has been one report that sperm whales ceased calling when exposed to pulses from a very distant seismic ship (Bowles *et al.*, 1994). However, more recent studies found that they continued calling in the presence of seismic pulses (Madsen *et al.*, 2002; Tyack *et al.*, 2003; Smultea *et al.*, 2004; Holst *et al.*, 2006; and Jochens *et al.*, 2008). Cerchio *et al.* (2014) suggested that the breeding display of humpback whales off Angola could have been disrupted by seismic sounds, as singing activity declined with increasing received levels. In addition, some cetaceans are known to change their calling rates, shift their peak frequencies, or otherwise modify their vocal behavior in response to airgun sounds (Di Iorio and Clark, 2010; Castellote *et al.*, 2012; Blackwell *et al.*, 2013). Di Iorio and Clark (2009) found evidence of increased calling by blue whales during operations by a lower-energy seismic source (*i.e.*, sparker). The hearing systems of baleen whales are undoubtedly more sensitive to low-frequency sounds than are the ears of small odontocetes that have been studied directly (MacGillivray *et al.*, 2013). Dolphins and porpoises commonly are heard calling while airguns are operating (*e.g.*, Gordon *et al.*, 2004; Smultea *et al.*, 2004; Holst *et al.*, 2005a, b; and Potter *et al.*, 2007). The sounds important to small odontocetes are predominantly at much higher frequencies than are the dominant components of airgun sounds, thus limiting the potential for masking.

Pinnipeds have the most sensitive hearing and/or produce most of their sounds in frequencies higher than the dominant components of airgun sound, but there is some overlap in the frequencies of the airgun pulses and the calls. However, the intermittent nature of airgun pulses presumably reduces the potential for masking.

Marine mammals are thought to be able to compensate for masking by adjusting their acoustic behavior through shifting call frequencies, increasing call volume, and increasing vocalization rates. For example blue whales are found to increase call rates when exposed to noise from seismic surveys in the St. Lawrence Estuary (Di

Iorio and Clark, 2009). The North Atlantic right whales (*Eubalaena glacialis*) exposed to high shipping noise increased call frequency (Parks *et al.*, 2007), while some humpback whales respond to low-frequency active sonar playbacks by increasing song length (Miller *et al.*, 2000). In general, NMFS expects the masking effects of seismic pulses to be minor, given the normally intermittent nature of seismic pulses.

Behavioral Disturbance

Marine mammals may behaviorally react to sound when exposed to anthropogenic noise. Disturbance includes a variety of effects, including subtle to conspicuous changes in behavior, movement, and displacement. Reactions to sound, if any, depend on species, state of maturity, experience, current activity, reproductive state, time of day, and many other factors (Richardson *et al.*, 1995; Wartzok *et al.*, 2004; Southall *et al.*, 2007; Weilgart, 2007; Ellison *et al.*, 2012). These behavioral reactions are often shown as: Changing durations of surfacing and dives, number of blows per surfacing, or moving direction and/or speed; reduced/increased vocal activities; changing/cessation of certain behavioral activities (such as socializing or feeding); visible startle response or aggressive behavior (such as tail/fluke slapping or jaw clapping); avoidance of areas where noise sources are located; and/or flight responses (*e.g.*, pinnipeds flushing into the water from haul-outs or rookeries). If a marine mammal does react briefly to an underwater sound by changing its behavior or moving a small distance, the impacts of the change are unlikely to be significant to the individual, let alone the stock or population (New *et al.*, 2013). However, if a sound source displaces marine mammals from an important feeding or breeding area for a prolonged period, impacts on individuals and populations could be significant (*e.g.*, Lusseau and Bejder, 2007; Weilgart, 2007).

The biological significance of many of these behavioral disturbances is difficult to predict, especially if the detected disturbances appear minor. However, the consequences of behavioral modification could be expected to be biologically significant if the change affects growth, survival, and/or reproduction. Some of these significant behavioral modifications include:

- Change in diving/surfacing patterns (such as those thought to be causing beaked whale stranding due to exposure to military mid-frequency tactical sonar);

- Habitat abandonment due to loss of desirable acoustic environment; and
- Cessation of feeding or social interaction.

The onset of behavioral disturbance from anthropogenic noise depends on both external factors (characteristics of noise sources and their paths) and the receiving animals (hearing, motivation, experience, demography) and is also difficult to predict (Richardson *et al.*, 1995; Southall *et al.*, 2007). Given the many uncertainties in predicting the quantity and types of impacts of noise on marine mammals, it is common practice to estimate how many mammals would be present within a particular distance of industrial activities and/or exposed to a particular level of sound. In most cases, this approach likely overestimates the numbers of marine mammals that would be affected in some biologically-important manner.

Baleen Whales—Baleen whales generally tend to avoid operating airguns, but avoidance radii are quite variable (reviewed in Richardson *et al.*, 1995; Gordon *et al.*, 2004). Whales are often reported to show no overt reactions to pulses from large arrays of airguns at distances beyond a few kilometers, even though the airgun pulses remain well above ambient noise levels out to much longer distances. However, baleen whales exposed to strong noise pulses from airguns often react by deviating from their normal migration route and/or interrupting their feeding and moving away. In the cases of migrating gray (*Eschrichtius robustus*) and bowhead (*Balaena mysticetus*) whales, the observed changes in behavior appeared to be of little or no biological consequence to the animals (Richardson *et al.*, 1995). They simply avoided the sound source by displacing their migration route to varying degrees, but within the natural boundaries of the migration corridors (Malme *et al.*, 1984; Malme and Miles, 1985; Richardson *et al.*, 1995).

Studies of gray, bowhead, and humpback whales have shown that seismic pulses with received levels of 160 to 170 dB re 1 μ Pa (rms) seem to cause obvious avoidance behavior in a substantial fraction of the animals exposed (Malme *et al.*, 1986, 1988; Richardson *et al.*, 1995). In many areas, seismic pulses from large arrays of airguns diminish to those levels at distances ranging from 4 to 15 km (2.2 to 8.1 nmi) from the source. A substantial proportion of the baleen whales within those distances may show avoidance or other strong behavioral reactions to the airgun array. Subtle behavioral changes sometimes

become evident at somewhat lower received levels, and studies have shown that some species of baleen whales, notably bowhead, gray, and humpback whales, at times, show strong avoidance at received levels lower than 160 to 170 dB re 1 μ Pa (rms).

Researchers have studied the responses of humpback whales to seismic surveys during migration, feeding during the summer months, breeding while offshore from Angola, and wintering offshore from Brazil. McCauley *et al.* (1998, 2000a) studied the responses of humpback whales off western Australia to a full-scale seismic survey with a 16 airgun array (2,678 in³) and to a single airgun (20 in³) with source level of 227 dB re 1 μ Pa (p-p). In the 1998 study, they documented that avoidance reactions began at 5 to 8 km (2.7 to 4.3 nmi) from the array, and that those reactions kept most pods approximately 3 to 4 km (1.6 to 2.2 nmi) from the operating seismic boat. In the 2000 study, they noted localized displacement during migration of 4 to 5 km (2.2 to 2.7 nmi) by traveling pods and 7 to 12 km (3.8 to 6.5 nmi) by more sensitive resting pods of cow-calf pairs. Avoidance distances with respect to the single airgun were smaller but consistent with the results from the full array in terms of the received sound levels. The mean received level for initial avoidance of an approaching airgun was 140 dB re 1 μ Pa (rms) for humpback pods containing females, and at the mean closest point of approach distance the received level was 143 dB re 1 μ Pa (rms). The initial avoidance response generally occurred at distances of 5 to 8 km (2.7 to 4.3 nmi) from the airgun array and 2 km (1.1 nmi) from the single airgun. However, some individual humpback whales, especially males, approached within distances of 100 to 400 m (328 to 1,312 ft), where the maximum received level was 179 dB re 1 μ Pa (rms). Studies examining the behavioral responses of humpback whales to airguns are currently underway off eastern Australia (Cato *et al.*, 2011, 2012, 2013).

Data collected by observers during several seismic surveys in the Northwest Atlantic showed that sighting rates of humpback whales were significantly greater during non-seismic periods compared with periods when a full array was operating (Moulton and Holst, 2010). In addition, humpback whales were more likely to swim away and less likely to swim towards a vessel during seismic vs. non-seismic periods (Moulton and Holst, 2010).

Humpback whales on their summer feeding grounds in southeast Alaska did not exhibit persistent avoidance when

exposed to seismic pulses from a 1.64–L (100 in³) airgun (Malme *et al.*, 1985). Some humpbacks seemed “startled” at received levels of 150 to 169 dB re 1 μ Pa. Malme *et al.* (1985) concluded that there was no clear evidence of avoidance, despite the possibility of subtle effects, at received levels up to 172 dB re 1 μ Pa (rms). However, Moulton and Holst (2010) reported that humpback whales monitored during seismic surveys in the Northwest Atlantic had lower sighting rates and were most often seen swimming away from the vessel during seismic periods compared with periods when airguns were silent.

Studies have suggested that South Atlantic humpback whales wintering off Brazil may be displaced or even strand upon exposure to seismic surveys (Engel *et al.*, 2004). The evidence for this was circumstantial and subject to alternative explanations (IAGC, 2004). Also, the evidence was not consistent with subsequent results from the same area of Brazil (Parente *et al.*, 2006), or with direct studies of humpbacks exposed to seismic surveys in other areas and seasons. After allowance for data from subsequent years, there was “no observable direct correlation” between strandings and seismic surveys (IWC, 2007: 236).

There are no reactions of right whales to seismic surveys. However, Rolland *et al.* (2012) suggested that ship noise causes increased stress in right whales; they showed that baseline levels of stress-related fecal hormone metabolites decreased in North Atlantic right whales with a 6 dB decrease in underwater noise from vessels. Wright *et al.* (2011) also reported that sound could be a potential source of stress for marine mammals.

Results from bowhead whales show that their responsiveness can be quite variable depending on their activity (migrating versus feeding). Bowhead whales migrating west across the Alaskan Beaufort Sea in autumn, in particular, are unusually responsive, with substantial avoidance occurring out to distances of 20 to 30 km (10.8 to 16.2 nmi) from a medium-sized airgun source (Miller *et al.*, 1999; Richardson *et al.*, 1999). However, more recent research on bowhead whales corroborates earlier evidence that, during the summer feeding season, bowheads are not as sensitive to seismic sources (Miller *et al.*, 2005). Nonetheless, Robertson *et al.* (2013) showed that bowheads on their summer feeding grounds showed subtle but statistically significant changes in surfacing-respiration-dive cycles during exposure to seismic sounds, including

shorter surfacing intervals, shorter dives, and decreased number of blows per surface interval.

Bowhead whale calls detected in the presence and absence of airgun sounds have been studied extensively in the Beaufort Sea. Bowheads continue to produce calls of the usual types when exposed to airgun sounds on their summering grounds, although number of calls detected are significantly lower in the presence than in the absence of airgun pulses; Blackwell *et al.* (2013) reported that calling rates in 2007 declined significantly where received SPLs from airgun sounds were 116 to 129 dB re 1 μ Pa. Thus, bowhead whales in the Beaufort Sea apparently decrease their calling rates in response to seismic operations, although movement out of the area could also contribute to the lower call detection rate (Blackwell *et al.*, 2013).

A multivariate analysis of factors affecting the distribution of calling bowhead whales during their fall migration in 2009 noted that the southern edge of the distribution of calling whales was significantly closer to shore with increasing levels of airgun sound from a seismic survey a few hundred kms to the east of the study area (*i.e.*, behind the westward-migrating whales; McDonald *et al.*, 2010, 2011). It was not known whether this statistical effect represented a stronger tendency for quieting of the whales farther offshore in deeper water upon exposure to airgun sound, or an actual inshore displacement of whales.

Reactions of migrating and feeding (but not wintering) gray whales to seismic surveys have been studied. Malme *et al.* (1986, 1988) studied the responses of feeding eastern Pacific gray whales to pulses from a single 100 in³ airgun off St. Lawrence Island in the northern Bering Sea. They estimated, based on small sample sizes, that 50 percent of feeding gray whales stopped feeding at an average received pressure level of 173 dB re 1 μ Pa on an (approximate) rms basis, and that 10 percent of feeding whales interrupted feeding at received levels of 163 dB re 1 μ Pa (rms). Those findings were generally consistent with the results of experiments conducted on larger numbers of gray whales that were migrating along the California coast (Malme *et al.*, 1984; Malme and Miles, 1985), and western Pacific gray whales feeding off Sakhalin Island, Russia (Wursig *et al.*, 1999; Gailey *et al.*, 2007; Johnson *et al.*, 2007; Yazvenko *et al.*, 2007a, b), along with data on gray whales off British Columbia (Bain and Williams, 2006).

Various species of *Balaenoptera* (blue, sei, fin, and minke whales) have occasionally been seen in areas ensounded by airgun pulses (Stone, 2003; MacLean and Haley, 2004; Stone and Tasker, 2006), and calls from blue and fin whales have been localized in areas with airgun operations (*e.g.*, McDonald *et al.*, 1995; Dunn and Hernandez, 2009; Castellote *et al.*, 2010). Sightings by observers on seismic vessels off the United Kingdom from 1997 to 2000 suggest that, during times of good sightability, sighting rates for mysticetes (mainly fin and sei whales) were similar when large arrays of airguns were shooting versus silent (Stone, 2003; Stone and Tasker, 2006). However, these whales tended to exhibit localized avoidance, remaining significantly further (on average) from the airgun array during seismic operations compared with non-seismic periods (Stone and Tasker, 2006). Castellote *et al.* (2010, 2012) reported that singing fin whales in the Mediterranean moved away from an operating airgun array, and their song notes had low bandwidths during periods with versus without airgun sounds.

Ship-based monitoring studies of baleen whales (including blue, fin, sei, minke, and humpback whales) in the Northwest Atlantic found that overall, this group had lower sighting rates during seismic vs. non-seismic periods (Moulton and Holst, 2010). Baleen whales as a group were also seen significantly farther from the vessel during seismic compared with non-seismic periods, and they were more often seen to be swimming away from the operating seismic vessel (Moulton and Holst, 2010). Blue and minke whales were initially sighted significantly farther from the vessel during seismic operations compared to non-seismic periods; the same trend was observed for fin whales (Moulton and Holst, 2010). Minke whales were most often observed to be swimming away from the vessel when seismic operations were underway (Moulton and Holst, 2010).

Data on short-term reactions by cetaceans to impulsive noises are not necessarily indicative of long-term or biologically significant effects. It is not known whether impulsive sounds affect reproductive rate or distribution and habitat use in subsequent days or years. However, gray whales have continued to migrate annually along the west coast of North America with substantial increases in the population over recent years, despite intermittent seismic exploration (and much ship traffic) in that area for decades (Appendix A in

Malme *et al.*, 1984; Richardson *et al.*, 1995; Allen and Angliss, 2010). The western Pacific gray whale population did not seem affected by a seismic survey in its feeding ground during a previous year (Johnson *et al.*, 2007). Similarly, bowhead whales have continued to travel to the eastern Beaufort Sea each summer, and their numbers have increased notably, despite seismic exploration in their summer and autumn range for many years (Richardson *et al.*, 1987; Allen and Angliss, 2010).

Toothed Whales—Little systematic information is available about reactions of toothed whales to noise pulses. Few studies similar to the more extensive baleen whale/seismic pulse work summarized above have been reported for toothed whales. However, there are recent systematic studies on sperm whales (*e.g.*, Gordon *et al.*, 2006; Madsen *et al.*, 2006; Winsor and Mate, 2006; Jochens *et al.*, 2008; Miller *et al.*, 2009). There is an increasing amount of information about responses of various odontocetes to seismic surveys based on monitoring studies (*e.g.*, Stone, 2003; Smultea *et al.*, 2004; Moulton and Miller, 2005; Bain and Williams, 2006; Holst *et al.*, 2006; Stone and Tasker, 2006; Potter *et al.*, 2007; Hauser *et al.*, 2008; Holst and Smultea, 2008; Weir, 2008; Barkaszi *et al.*, 2009; Richardson *et al.*, 2009; Moulton and Holst, 2010).

Seismic operators and PSOs on seismic vessels regularly see dolphins and other small toothed whales near operating airgun arrays, but in general there is a tendency for most delphinids to show some avoidance of operating seismic vessels (*e.g.*, Goold, 1996a,b,c; Calambokidis and Osmeck, 1998; Stone, 2003; Moulton and Miller, 2005; Holst *et al.*, 2006; Stone and Tasker, 2006; Weir, 2008; Richardson *et al.*, 2009; Barkaszi *et al.*, 2009; Moulton and Holst, 2010; Barry *et al.*, 2012). Some dolphins seem to be attracted to the seismic vessel and floats, and some ride the bow wave of the seismic vessel even when large arrays of airguns are firing (*e.g.*, Moulton and Miller, 2005). Nonetheless, small toothed whales more often tend to head away, or to maintain a somewhat greater distance from the vessel, when a large array of airguns is operating than when it is silent (*e.g.*, Stone and Tasker, 2006; Weir, 2008; Barry *et al.*, 2010; Moulton and Holst, 2010). In most cases, the avoidance radii for delphinids appear to be small, on the order of one km or less, and some individuals show no apparent avoidance. Captive bottlenose dolphins (*Tursiops truncatus*) and beluga whales (*Delphinapterus leucas*) exhibited changes in behavior when exposed to

strong pulsed sounds similar in duration to those typically used in seismic surveys (Finneran *et al.*, 2000, 2002, 2005). However, the animals tolerated high received levels of sound before exhibiting aversive behaviors.

Preliminary findings of a monitoring study of narwhals (*Monodon monoceros*) in Melville Bay, Greenland (summer and fall 2012) showed no short-term effects of seismic survey activity on narwhal distribution, abundance, migration timing, and feeding habits (Heide-Jorgensen *et al.*, 2013a). In addition, there were no reported effects on narwhal hunting. These findings do not seemingly support a suggestion by Heide-Jorgensen *et al.* (2013b) that seismic surveys in Baffin Bay may have delayed the migration timing of narwhals, thereby increasing the risk of narwhals to ice entrapment.

Results of porpoises depend on species. The limited available data suggest that harbor porpoises (*Phocoena phocoena*) show stronger avoidance of seismic operations than do Dall's porpoises (*Phocoenoides dalli*) (Stone, 2003; MacLean and Koski, 2005; Bain and Williams, 2006; Stone and Tasker, 2006). Thompson *et al.* (2013) reported decreased densities and reduced acoustic detections of harbor porpoise in response to a seismic survey in Moray Firth, Scotland, at ranges of 5 to 10 km (2.7 to 5.4 nmi) (SPLs of 165 to 172 dB re 1 μ Pa; sound exposure levels (SELs) of 145 to 151 dB μ Pa²s); however, animals returned to the area within a few hours. Dall's porpoises seem relatively tolerant of airgun operations (MacLean and Koski, 2005; Bain and Williams, 2006), although they too have been observed to avoid large arrays of operating airguns (Calambokidis and Osmeck, 1998; Bain and Williams, 2006). This apparent difference in responsiveness of these two porpoise species is consistent with their relative responsiveness to boat traffic and some other acoustic sources (Richardson *et al.*, 1995; Southall *et al.*, 2007).

Most studies of sperm whales exposed to airgun sounds indicate that the sperm whale shows considerable tolerance of airgun pulses (*e.g.*, Stone, 2003; Moulton *et al.*, 2005, 2006a; Stone and Tasker, 2006; Weir, 2008). In most cases the whales do not show strong avoidance, and they continue to call. However, controlled exposure experiments in the Gulf of Mexico indicate that foraging behavior was altered upon exposure to airgun sound (Jochens *et al.*, 2008; Miller *et al.*, 2009; Tyack, 2009). There are almost no specific data on the behavioral reactions of beaked whales to seismic surveys.

However, some northern bottlenose whales (*Hyperoodon ampullatus*) remained in the general area and continued to produce high-frequency clicks when exposed to sound pulses from distant seismic surveys (Gosselin and Lawson, 2004; Laurinolli and Cochrane, 2005; Simard *et al.*, 2005). Most beaked whales tend to avoid approaching vessels of other types (*e.g.*, Wursig *et al.*, 1998). They may also dive for an extended period when approached by a vessel (*e.g.*, Kasuya, 1986), although it is uncertain how much longer such dives may be as compared to dives by undisturbed beaked whales, which also are often quite long (Baird *et al.*, 2006; Tyack *et al.*, 2006). Based on a single observation, Aguilar-Soto *et al.* (2006) suggested that foraging efficiency of Cuvier's beaked whales may be reduced by close approach of vessels. In any event, it is likely that most beaked whales would also show strong avoidance of an approaching seismic vessel, although this has not been documented explicitly. In fact, Moulton and Holst (2010) reported 15 sightings of beaked whales during seismic studies in the Northwest Atlantic; seven of those sightings were made at times when at least one airgun was operating. There was little evidence to indicate that beaked whale behavior was affected by airgun operations; sighting rates and distances were similar during seismic and non-seismic periods (Moulton and Holst, 2010).

There are increasing indications that some beaked whales tend to strand when naval exercises involving mid-frequency sonar operation are ongoing nearby (*e.g.*, Simmonds and Lopez-Jurado, 1991; Frantzis, 1998; NOAA and USN, 2001; Jepson *et al.*, 2003; Hildebrand, 2005; Barlow and Gisiner, 2006; see also the "Stranding and Mortality" section in this notice). These strandings are apparently a disturbance response, although auditory or other injuries or other physiological effects may also be involved. Whether beaked whales would ever react similarly to seismic surveys is unknown. Seismic survey sounds are quite different from those of the sonar in operation during the above-cited incidents.

Odontocete reactions to large arrays of airguns are variable and, at least for delphinids, seem to be confined to a smaller radius than has been observed for the more responsive of some mysticetes. However, other data suggest that some odontocete species, including harbor porpoises, may be more responsive than might be expected given their poor low-frequency hearing. Reactions at longer distances may be

particularly likely when sound propagation conditions are conducive to transmission of the higher frequency components of airgun sound to the animals' location (DeRuiter *et al.*, 2006; Goold and Coates, 2006; Tyack *et al.*, 2006; Potter *et al.*, 2007).

Pinnipeds—Pinnipeds are not likely to show a strong avoidance reaction to the airgun array. Visual monitoring from seismic vessels has shown only slight (if any) avoidance of airguns by pinnipeds, and only slight (if any) changes in behavior. In the Beaufort Sea, some ringed seals avoided an area of 100 m to (at most) a few hundred meters around seismic vessels, but many seals remained within 100 to 200 m (328 to 656 ft) of the trackline as the operating airgun array passed by (*e.g.*, Harris *et al.*, 2001; Moulton and Lawson, 2002; Miller *et al.*, 2005.). Ringed seal (*Pusa hispida*) sightings averaged somewhat farther away from the seismic vessel when the airguns were operating than when they were not, but the difference was small (Moulton and Lawson, 2002). Similarly, in Puget Sound, sighting distances for harbor seals (*Phoca vitulina*) and California sea lions (*Zalophus californianus*) tended to be larger when airguns were operating (Calambokidis and Osmeck, 1998). Previous telemetry work suggests that avoidance and other behavioral reactions by two other species of seals to small airgun sources may be stronger than evident to date from visual studies of pinnipeds reactions to airguns (Thompson *et al.*, 1998).

During seismic exploration off Nova Scotia, gray seals (*Halichoerus grypus*) exposed to noise from airguns and linear explosive charges did not react strongly (J. Parsons in Greene *et al.*, 1985). Pinnipeds in both water and air, sometimes tolerate strong noise pulses from non-explosive and explosive scaring devices, especially if attracted to the area for feeding and reproduction (Mate and Harvey, 1987; Reeves *et al.*, 1996). Thus pinnipeds are expected to be rather tolerant of, or habituate to, repeated underwater sounds from distant seismic sources, at least when the animals are strongly attracted to the area.

Hearing Impairment and Other Physical Effects

Exposure to high intensity sound for a sufficient duration may result in auditory effects such as a noise-induced threshold shift—an increase in the auditory threshold after exposure to noise (Finneran, Carder, Schlundt, and Ridgway, 2005). Factors that influence the amount of threshold shift include the amplitude, duration, frequency

content, temporal pattern, and energy distribution of noise exposure. The magnitude of hearing threshold shift normally decreases over time following cessation of the noise exposure. The amount of threshold shift just after exposure is called the initial threshold shift. If the threshold shift eventually returns to zero (*i.e.*, the threshold returns to the pre-exposure value), it is called temporary threshold shift (TTS) (Southall *et al.*, 2007). Researchers have studied TTS in certain captive odontocetes and pinnipeds exposed to strong sounds (reviewed in Southall *et al.*, 2007). However, there has been no specific documentation of TTS, let alone permanent hearing damage, *i.e.*, permanent threshold shift (PTS), in free-ranging marine mammals exposed to sequences of airgun pulses during realistic field conditions.

Temporary Threshold Shift—TTS is the mildest form of hearing impairment that can occur during exposure to a strong sound (Kryter, 1985). While experiencing TTS, the hearing threshold rises and a sound must be stronger in order to be heard. At least in terrestrial mammals, TTS can last from minutes or hours to (in cases of strong TTS) days. For sound exposures at or somewhat above the TTS threshold, hearing sensitivity in both terrestrial and marine mammals recovers rapidly after exposure to the noise ends. Few data on sound levels and durations necessary to elicit mild TTS have been obtained for marine mammals, and none of the published data concern TTS elicited by exposure to multiple pulses of sound. Available data on TTS in marine mammals are summarized in Southall *et al.* (2007). Table 2 (above) presents the estimated distances from the *Revelle's* airguns at which the received energy level (per pulse, flat-weighted) would be expected to be greater than or equal to 180 and 190 dB re 1 μ Pa (rms).

The established 180 and 190 dB (rms) criteria are not considered to be the levels above which TTS might occur. Rather, they are the received levels above which, in the view of a panel of bioacoustics specialists convened by NMFS before TTS measurements for marine mammals started to become available, one could not be certain that there would be no injurious effects, auditory or otherwise, to marine mammals. NMFS also assumes that cetaceans and pinnipeds exposed to levels exceeding 160 dB re 1 μ Pa (rms) may experience Level B harassment.

For toothed whales, researchers have derived TTS information for odontocetes from studies on the bottlenose dolphin and beluga. The experiments show that exposure to a

single impulse at a received level of 207 kPa (or 30 psi, peak-to-peak), which is equivalent to 228 dB re 1 Pa (peak-to-peak), resulted in a 7 and 6 dB TTS in the beluga whale at 0.4 and 30 kHz, respectively. Thresholds returned to within 2 dB of the pre-exposure level within 4 minutes of the exposure (Finneran *et al.*, 2002). For the one harbor porpoise tested, the received level of airgun sound that elicited onset of TTS was lower (Lucke *et al.*, 2009). If these results from a single animal are representative, it is inappropriate to assume that onset of TTS occurs at similar received levels in all odontocetes (*cf.* Southall *et al.*, 2007). Some cetaceans apparently can incur TTS at considerably lower sound exposures than are necessary to elicit TTS in the beluga or bottlenose dolphin.

For baleen whales, there are no data, direct or indirect, on levels or properties of sound that are required to induce TTS. The frequencies to which baleen whales are most sensitive are assumed to be lower than those to which odontocetes are most sensitive, and natural background noise levels at those low frequencies tend to be higher. As a result, auditory thresholds of baleen whales within their frequency band of best hearing are believed to be higher (less sensitive) than are those of odontocetes at their best frequencies (Clark and Ellison, 2004). From this, it is suspected that received levels causing TTS onset may also be higher in baleen whales than those of odontocetes (Southall *et al.*, 2007).

In pinnipeds, researchers have not measured TTS thresholds associated with exposure to brief pulses (single or multiple) of underwater sound. Initial evidence from more prolonged (non-pulse) exposures suggested that some pinnipeds (harbor seals in particular) incur TTS at somewhat lower received levels than do small odontocetes exposed for similar durations (Kastak *et al.*, 1999, 2005; Ketten *et al.*, 2001). The TTS threshold for pulsed sounds has been indirectly estimated as being an SEL of approximately 171 dB re 1 μ Pa²-s (Southall *et al.*, 2007) which would be equivalent to a single pulse with a received level of approximately 181 to 186 dB re 1 μ Pa (rms), or a series of pulses for which the highest rms values are a few dB lower. Corresponding values for California sea lions and northern elephant seals (*Mirounga angustirostris*) are likely to be higher (Kastak *et al.*, 2005).

Additional data are needed to determine the received levels at which small odontocetes would start to incur TTS upon exposure to repeated, low-frequency pulses of airgun sounds with

variable received levels. To determine how close an airgun array would need to approach in order to elicit TTS, one would (as a minimum) need to allow for the sequence of distances at which airgun pulses would occur, and for the dependence of received SEL on distance in the region of the airgun operation (Breitzke and Bohlen, 2010; Laws, 2012). At the present state of knowledge, it can be assumed that the effect is directly related to total received energy, although there is recent evidence that auditory effects in a given animal are not a simple function of received acoustic energy. Frequency, duration of the exposure and occurrence of gaps within the exposure can also influence the auditory effect (Finneran and Schlundt, 2010, 2011, 2013; Finneran *et al.*, 2010a,b; Finneran 2012; Ketten, 2012; Kastelein *et al.*, 2013a).

The assumption that, in marine mammals, the occurrence and magnitude of TTS is a function of cumulative acoustic energy (SEL) is probably an oversimplification (Finneran, 2012). Popov *et al.* (2011) examined the effects of fatiguing noise on the hearing threshold of Yangtze finless porpoises (*Neophocaena phocaenoides*) when exposed to frequencies of 32 to 128 kHz at 140 to 160 dB re 1 μ Pa for 1 to 30 minutes. They found that an exposure of higher level and shorter duration produced a higher TTS than an exposure of equal SEL but of lower level and longer duration. Kastelein *et al.* (2012a,b; 2013b) also reported that the equal-energy model is not valid for predicting TTS in harbor porpoises or harbor seals.

Recent data have shown that the SEL required for TTS onset to occur increases with intermittent exposures, with some auditory recovery during silent periods between (Finneran *et al.*, 2010b; Finneran and Schlundt, 2011). Schlundt *et al.* (2013) reported that the potential for seismic surveys using airguns to cause auditory effects on dolphins could be lower than previously thought. Based on behavioral tests, Finneran *et al.* (2011) and Schlundt *et al.* (2013) reported no measurable TTS in bottlenose dolphins after exposure to 10 impulses from a seismic airgun with a cumulative SEL of approximately 195 dB re 1 μ Pa²-s; results from auditory evoked potential measurements were more variable (Schlundt *et al.*, 2013).

Recent studies have also shown that the SEL necessary to elicit TTS can depend substantially on frequency, with susceptibility to TTS increasing with increasing frequency above 3 kHz (Finneran and Schlundt, 2010, 2011; Finneran, 2012). When beluga whales

were exposed to fatiguing noise with sound levels of 165 dB re 1 μ Pa for durations of 1 to 30 minutes at frequencies of 11.2 to 90 kHz, the highest TTS with the longest recovery time was produced by lower frequencies (11.2 and 22.5 kHz); TTS effects also gradually increased with prolonged exposure time (Popov *et al.*, 2013a). Popov *et al.* (2013b) also reported that TTS produced by exposure to a fatiguing noise was larger during the first session (or naïve subject state) with a beluga whale than TTS that resulted from the same sound in subsequent sessions (experienced subject state). Therefore, Supin *et al.* (2013) reported that SEL may not be a valid metric for examining fatiguing sounds on beluga whales. Similarly, Nachtigall and Supin (2013) reported that false killer whales are able to change their hearing sensation levels when exposed to loud sounds, such as warning signals or echolocation sounds.

It is inappropriate to assume that onset of TTS occurs at similar received levels in all cetaceans (Southall *et al.*, 2007). Some cetaceans could incur TTS at lower sound exposures than are necessary to elicit TTS in the beluga or bottlenose dolphin. Based on the best available information, Southall *et al.* (2007) recommended a TTS threshold for exposure to a single or multiple pulses of 183 dB re 1 μ Pa²s. Tougaard *et al.* (2013) proposed a TTS criterion of 165 dB re 1 μ Pa²s for porpoises based on data from two recent studies. Gedamke *et al.* (2011), based on preliminary simulation modeling that attempted to allow for various uncertainties in assumptions and variability around population means, suggested that some baleen whales whose closest point of approach to a seismic vessel is 1 km or more could experience TTS.

Permanent Threshold Shift—When PTS occurs, there is physical damage to the sound receptors in the ear. In severe cases, there can be total or partial deafness, whereas in other cases, the animal has an impaired ability to hear sounds in specific frequency ranges (Kryter, 1985). There is no specific evidence that exposure to pulses of airgun sound can cause PTS in any marine mammal, even with large arrays of airguns. However, given the possibility that mammals close to an airgun array might incur at least mild TTS, there has been further speculation about the possibility that some individuals occurring very close to airguns might incur PTS (*e.g.*, Richardson *et al.*, 1995, p. 372ff; Gedamke *et al.*, 2008). Single or occasional occurrences of mild TTS are

not indicative of permanent auditory damage, but repeated or (in some cases) single exposures to a level well above that causing TTS onset might elicit PTS.

Relationships between TTS and PTS thresholds have not been studied in marine mammals but are assumed to be similar to those in humans and other terrestrial mammals (Southall *et al.*, 2007). PTS might occur at a received sound level at least several dBs above that inducing mild TTS if the animal were exposed to strong sound pulses with rapid rise times. Based on data from terrestrial mammals, a precautionary assumption is that the PTS threshold for impulse sounds (such as airgun pulses as received close to the source) is at least 6 dB higher than the TTS threshold on a peak-pressure basis, and probably greater than 6 dB (Southall *et al.*, 2007). Given the higher level of sound necessary to cause PTS as compared with TTS, it is considerably less likely that PTS would occur. Baleen whales generally avoid the immediate area around operating seismic vessels, as do some other marine mammals.

Non-auditory Physiological Effects—Non-auditory physiological effects or injuries that theoretically might occur in marine mammals exposed to strong underwater sound include stress, neurological effects, bubble formation, resonance, and other types of organ or tissue damage (Cox *et al.*, 2006; Southall *et al.*, 2007). Studies examining such effects are limited. However, resonance effects (Gentry, 2002) and direct noise-induced bubble formations (Crum *et al.*, 2005) are implausible in the case of exposure to an impulsive broadband source like an airgun array. If seismic surveys disrupt diving patterns of deep-diving species, this might perhaps result in bubble formation and a form of the bends, as speculated to occur in beaked whales exposed to sonar. However, there is no specific evidence of this upon exposure to airgun pulses.

In general, very little is known about the potential for seismic survey sounds (or other types of strong underwater sounds) to cause non-auditory physical effects in marine mammals. Such effects, if they occur at all, would presumably be limited to short distances and to activities that extend over a prolonged period. The available data do not allow identification of a specific exposure level above which non-auditory effects can be expected (Southall *et al.*, 2007), or any meaningful quantitative predictions of the numbers (if any) of marine mammals that might be affected in those ways. Marine mammals that show behavioral avoidance of seismic vessels, including most baleen whales, some odontocetes,

and some pinnipeds, are especially unlikely to incur non-auditory physical effects.

There is no definitive evidence that any of these effects occur even for marine mammals in close proximity to large airgun arrays. However, Gray and Van Waerebeek (2011) have suggested a cause-effect relationship between a seismic survey off Liberia in 2009 and the erratic movement, postural instability, and akinesia in a pantropical spotted dolphin based on spatially and temporally close association with the airgun array. Additionally, a few cases of strandings in the general area where a seismic survey was ongoing have led to speculation concerning a possible link between seismic surveys and strandings (Castellote and Llorens, 2013).

Stranding and Mortality—When a living or dead marine mammal swims or floats onto shore and becomes “beached” or incapable of returning to sea, the event is termed a “stranding” (Geraci *et al.*, 1999; Perrin and Geraci, 2002; Geraci and Lounsbury, 2005; NMFS, 2007). The legal definition for a stranding under the MMPA is that “(A) a marine mammal is dead and is (i) on a beach or shore of the United States; or (ii) in waters under the jurisdiction of the United States (including any navigable waters); or (B) a marine mammal is alive and is (i) on a beach or shore of the United States and is unable to return to the water; (ii) on a beach or shore of the United States and, although able to return to the water is in need of apparent medical attention; or (iii) in the waters under the jurisdiction of the United States (including any navigable waters), but is unable to return to its natural habitat under its own power or without assistance.”

Marine mammals are known to strand for a variety of reasons, such as infectious agents, biotoxins, starvation, fishery interaction, ship strike, unusual oceanographic or weather events, sound exposure, or combinations of these stressors sustained concurrently or in series. However, the cause or causes of most strandings are unknown (Geraci *et al.*, 1976; Eaton, 1979; Odell *et al.*, 1980; Best, 1982). Numerous studies suggest that the physiology, behavior, habitat relationships, age, or condition of cetaceans may cause them to strand or might pre-dispose them to strand when exposed to another phenomenon. These suggestions are consistent with the conclusions of numerous other studies that have demonstrated that combinations of dissimilar stressors commonly combine to kill an animal or

dramatically reduce its fitness, even though one exposure without the other does not produce the same result (Chroussos, 2000; Creel, 2005; DeVries *et al.*, 2003; Fair and Becker, 2000; Foley *et al.*, 2001; Moberg, 2000; Relyea, 2005a, 2005b; Romero, 2004; Sih *et al.*, 2004).

Strandings Associated With Military Active Sonar—The proposed action is not a military readiness activity or using military active sonar (non-pulse). Several sources have published lists of mass stranding events of cetaceans in an attempt to identify relationships between those stranding events and military active sonar (Hildebrand, 2004; IWC, 2005; Taylor *et al.*, 2004). For example, based on a review of stranding records between 1960 and 1995, the International Whaling Commission (2005) identified ten mass stranding events and concluded that, out of eight stranding events reported from the mid-1980s to the summer of 2003, seven had been coincident with the use of mid-frequency active sonar and most involved beaked whales.

Over the past 12 years, there have been five stranding events coincident with military mid-frequency active sonar use in which exposure to sonar is believed to have been a contributing factor to strandings: Greece (1996); the Bahamas (2000); Madeira (2000); Canary Islands (2002); and Spain (2006). Refer to Cox *et al.* (2006) for a summary of common features shared by the strandings events in Greece (1996), Bahamas (2000), Madeira (2000), and Canary Islands (2002); and Fernandez *et al.*, (2005) for an additional summary of the Canary Islands 2002 stranding event.

Potential for Stranding From Seismic Surveys—Marine mammals close to underwater detonations of high explosives can be killed or severely injured, and the auditory organs are especially susceptible to injury (Ketten *et al.*, 1993; Ketten, 1995). However, explosives are no longer used in marine waters for commercial seismic surveys or (with rare exceptions) for seismic research. These methods have been replaced entirely by airguns or related non-explosive pulse generators. Airgun pulses are less energetic and have slower rise times, and there is no specific evidence that they can cause serious injury, death, or stranding even in the case of large airgun arrays. However, the association of strandings of beaked whales with naval exercises involving mid-frequency active sonar (non-pulse sound) and, in one case, the regional co-occurrence of an L-DEO seismic survey (Malakoff, 2002; Cox *et al.*, 2006), has raised the possibility that beaked whales exposed to strong

“pulsed” sounds could also be susceptible to injury and/or behavioral reactions that can lead to stranding (*e.g.*, Hildebrand, 2005; Southall *et al.*, 2007).

Specific sound-related processes that lead to strandings and mortality are not well documented, but may include:

- (1) Swimming in avoidance of a sound into shallow water;
- (2) A change in behavior (such as a change in diving behavior) that might contribute to tissue damage, gas bubble formation, hypoxia, cardiac arrhythmia, hypertensive hemorrhage or other forms of trauma;
- (3) A physiological change such as a vestibular response leading to a behavioral change or stress-induced hemorrhagic diathesis, leading in turn to tissue damage; and
- (4) Tissue damage directly from sound exposure, such as through acoustically-mediated bubble formation and growth or acoustic resonance of tissues.

Some of these mechanisms are unlikely to apply in the case of impulse sounds. However, there are indications that gas-bubble disease (analogous to “the bends”), induced in supersaturated tissue by a behavioral response to acoustic exposure, could be a pathologic mechanism for the strandings and mortality of some deep-diving cetaceans exposed to sonar. The evidence for this remains circumstantial and associated with exposure to naval mid-frequency sonar, not seismic surveys (Cox *et al.*, 2006; Southall *et al.*, 2007).

Seismic pulses and mid-frequency sonar signals are quite different, and some mechanisms by which sonar sounds have been hypothesized to affect beaked whales are unlikely to apply to airgun pulses. Sounds produced by airgun arrays are broadband impulses with most of the energy below one kHz. Typical military mid-frequency sonar emits non-impulse sounds at frequencies of 2 to 10 kHz, generally with a relatively narrow bandwidth at any one time. A further difference between seismic surveys and naval exercises is that naval exercises can involve sound sources on more than one vessel. Thus, it is not appropriate to expect that the same effects to marine mammals would result from military sonar and seismic surveys. However, evidence that sonar signals can, in special circumstances, lead (at least indirectly) to physical damage and mortality (*e.g.*, Balcomb and Claridge, 2001; NOAA and USN, 2001; Jepson *et al.*, 2003; Fernández *et al.*, 2004, 2005; Hildebrand 2005; Cox *et al.*, 2006) suggests that caution is warranted when dealing with exposure of marine mammals to any high-intensity sound.

There is no conclusive evidence of cetacean strandings or deaths at sea as a result of exposure to seismic surveys, but a few cases of strandings in the general area where a seismic survey was ongoing have led to speculation concerning a possible link between seismic surveys and strandings. Suggestions that there was a link between seismic surveys and strandings of humpback whales in Brazil (Engel *et al.*, 2004) were not well founded (IAGC, 2004; IWC, 2007). In September 2002, there was a stranding of two Cuvier's beaked whales in the Gulf of California, Mexico, when the L-DEO vessel R/V *Maurice Ewing* was operating a 20 airgun (8,490 in³) array in the general region. The link between the stranding and the seismic surveys was inconclusive and not based on any physical evidence (Hogarth, 2002; Yoder, 2002). Nonetheless, the Gulf of California incident plus the beaked whale strandings near naval exercises involving use of mid-frequency sonar suggests a need for caution in conducting seismic surveys in areas occupied by beaked whales until more is known about effects of seismic surveys on those species (Hildebrand, 2005). No injuries of beaked whales are anticipated during the proposed study because of:

- (1) The high likelihood that any beaked whales nearby would avoid the approaching vessel before being exposed to high sound levels, and
- (2) Differences between the sound sources to be used in the proposed study and operated by SIO and those involved in the naval exercises associated with strandings.

Potential Effects of Other Acoustic Devices and Sources

Multi-Beam Echosounder

SIO would operate the Kongsberg EM 122 multi-beam echosounder from the source vessel during the planned study. Sounds from the multi-beam echosounder are very short pulses, occurring for approximately 2 to 15 ms once every 5 to 20 seconds, depending on water depth. Most of the energy in the sound pulses emitted by the multi-beam echosounder is at frequencies near 12 kHz (10.5 to 13), and the maximum source level is 242 dB re 1 μ Pa (rms). The beam is narrow (1 to 2°) in fore-aft extent and wide (150°) in the cross-track extent. Each ping consists of eight (in water greater than 1,000 m deep) or four (in water less than 1,000 m) consecutive successive fan-shaped transmissions (segments) at different cross-track angles. Any given marine mammal at depth near the trackline would be in the

main beam for only one or two of the eight segments. Also, marine mammals that encounter the Kongsberg EM 122 are unlikely to be subjected to repeated pulses because of the narrow fore-aft width of the beam and would receive only limited amounts of pulse energy because of the short pulses. Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensonified for more than one 2 to 15 ms pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when a multi-beam echosounder emits a pulse is small. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans: (1) Generally have longer pulse duration than the Kongsberg EM 122; and (2) are often directed close to horizontally, as well as omnidirectional, versus more downward and narrowly for the multi-beam echosounder. The area of possible influence of the multi-beam echosounder is much smaller—a narrow band below the source vessel. Also, the duration of exposure for a given marine mammal can be much longer for naval sonar. During SIO's operations, the individual pulses would be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by. Possible effects of a multi-beam echosounder on marine mammals are described below.

Stranding—In 2013, an International Scientific Review Panel investigated a 2008 mass stranding of approximately 100 melon-headed whales in a Madagascar lagoon system (Southall *et al.*, 2013) associated with the use of a high-frequency mapping system. The report indicated that the use of a 12 kHz multi-beam echosounder was the most plausible and likely initial behavioral trigger of the mass stranding event. This was the first time that a relatively high-frequency mapping sonar system has been associated with a stranding event. However, the report also notes that there were several site- and situation-specific secondary factors that may have contributed to the avoidance responses that lead to the eventual entrapment and mortality of the whales within the Loza Lagoon system (*e.g.*, the survey vessel transiting in a north-south direction on the shelf break parallel to the shore may have trapped the animals between the sound source and the shore driving them towards the Loza Lagoon). The

report concluded that for odontocete cetaceans that hear well in the 10 to 50 kHz range, where ambient noise is typically quite low, high-power active sonars operating in this range may be more easily audible and have potential effects over larger areas than low-frequency systems that have more typically been considered in terms of anthropogenic noise impacts (Southall *et al.*, 2013). However, the risk may be very low given the extensive use of these systems worldwide on a daily basis and the lack of direct evidence of such responses previously (Southall *et al.*, 2013). It is noted that leading scientific experts on multi-beam echosounders have expressed concerns about the independent scientific review panel analyses and findings (Bernstein, 2013).

Masking—Marine mammal communications would not be masked appreciably by the multi-beam echosounder signals, given the low duty cycle of the echosounder and the brief period when an individual mammal is likely to be within its beam. Furthermore, in the case of baleen whales, the multi-beam echosounder signals (12 kHz) generally do not overlap with the predominant frequencies in the calls (16 Hz to less than 12 kHz), which would avoid any significant masking (Richardson *et al.*, 1995).

Behavioral Responses—Behavioral reactions of free-ranging marine mammals to sonars, echosounders, and other sound sources appear to vary by species and circumstance. Observed reactions have included silencing and dispersal by sperm whales (Watkins *et al.*, 1985), increased vocalizations and no dispersal by pilot whales (Rendell and Gordon, 1999), and the previously-mentioned beachings by beaked whales. During exposure to a 21 to 25 kHz “whale-finding” sonar with a source level of 215 dB re 1 μ Pa, gray whales reacted by orienting slightly away from the source and being deflected from their course by approximately 200 m (656.2 ft) (Frankel, 2005). When a 38 kHz echosounder and a 150 kHz ADCP were transmitting during studies in the Eastern Tropical Pacific, baleen whales showed no significant responses, while spotted and spinner dolphins were detected slightly more often and beaked whales less often during visual surveys (Gerrodette and Pettis, 2005).

Captive bottlenose dolphins and a beluga whale exhibited changes in behavior when exposed to 1 second tonal signals at frequencies similar to those that would be emitted by the multi-beam echosounder used by SIO, and to shorter broadband pulsed signals.

Behavioral changes typically involved what appeared to be deliberate attempts to avoid the sound exposure (Schlundt *et al.*, 2000; Finneran *et al.*, 2002; Finneran and Schlundt, 2004). The relevance of those data to free-ranging odontocetes is uncertain, and in any case, the test sounds were quite different in duration as compared with those from a multi-beam echosounder.

Risch *et al.* (2012) found a reduction in humpback whale song in the Stellwagen Bank National Marine Sanctuary during Ocean Acoustic Waveguide Remote Sensing (OAWRS) activities that were carried out approximately 200 km (108 nmi) away. The OAWRS used three frequency-modulated pulses centered at frequencies of 415, 734, and 949 Hz with received levels in the sanctuary of 88 to 110 dB re 1 μ Pa. Deng *et al.* (2014) measured the spectral properties of pulses transmitted by three 200 kHz echosounders, and found that they generated weaker sounds at frequencies below the center frequency (90 to 130 kHz). These sounds are within the hearing range of some marine mammals, and the authors suggested that they could be strong enough to elicit behavioral responses within close proximity to the sources, although they would be well below potentially harmful levels.

Hearing Impairment and Other Physical Effects—Given several stranding events that have been associated with the operation of naval sonar in specific circumstances, there is concern that mid-frequency sonar sounds can cause serious impacts to marine mammals (see above). However, the multi-beam echosounder proposed for use by SIO is quite different than sonar used for Navy operations. Pulse duration of the multi-beam echosounder is very short relative to the naval sonar. Also, at any given location, an individual marine mammal would be in the beam of the multi-beam echosounder for much less time, given the generally downward orientation of the beam and its narrow fore-aft beamwidth; Navy sonar often uses near-horizontally-directed sound and have higher duty cycles. Those factors would all reduce the sound energy received from the multi-beam echosounder rather drastically relative to that from naval sonar. NMFS believes that the brief exposure of marine mammals to one pulse, or small numbers of signals, from the multi-beam echosounder in this particular case is not likely to result in the harassment of marine mammals.

Sub-Bottom Profiler

SIO would operate a sub-bottom profiler (Knudsen 3260) from the source vessel during the proposed study. Sounds from the sub-bottom profiler are very short pulses, occurring for 1 to 4 ms once ever second. Most of the energy in the sound pulses emitted by the sub-bottom profiler is at frequencies 3.5 kHz, and the beam is directed downward. The sub-bottom profiler that may be used on the *Revelle* has a maximum source level of 204 dB re 1 μ Pa. The sonar emits energy in a 27° beam from the bottom of the ship. Marine mammals that encounter the Knudsen 3260 are unlikely to be subjected to repeated pulses because of the relatively narrow fore-aft width of the beam and would receive only limited amounts of pulse energy because of the short pulses. Animals close to the ship (where the beam is narrowest) are especially unlikely to be ensonified for more than one pulse (or two pulses if in the overlap area). Similarly, Kremser *et al.* (2005) noted that the probability of a cetacean swimming through the area of exposure when a sub-bottom profiler emits a pulse is small—even for a sub-bottom profiler more powerful than that that may be on the *Revelle*. The animal would have to pass the transducer at close range and be swimming at speeds similar to the vessel in order to receive the multiple pulses that might result in sufficient exposure to cause TTS.

Navy sonars that have been linked to avoidance reactions and stranding of cetaceans: (1) Generally have longer pulse duration than the Knudsen 3260; and (2) are often directed close to horizontally versus more downward for the sub-bottom profiler. The area of possible influence of the single-beam echosounder is much smaller—a narrow band below the source vessel. Also, the duration of exposure for a given marine mammal can be much longer for naval sonar. During SIO's operations, the individual pulses would be very short, and a given mammal would not receive many of the downward-directed pulses as the vessel passes by. Possible effects of a sub-bottom profiler on marine mammals are described below.

Masking—Marine mammal communications would not be masked appreciably by the sub-bottom profiler signals given the directionality of the signal and the brief period when an individual mammal is likely to be within its beam. Furthermore, in the case of baleen whales, the sub-bottom profiler signals do not overlap with the predominant frequencies in the calls (16 Hz to less than 12 kHz), which would

avoid any significant masking (Richardson *et al.*, 1995).

Behavioral Responses—Marine mammal behavioral reactions to other pulsed sound sources are discussed above, and responses to the sub-bottom profiler are likely to be similar to those for other pulsed sources if received at the same levels. However, the pulsed signals from the sub-bottom profiler are considerably weaker than those from the multi-beam echosounder. Therefore, behavioral responses are not expected unless marine mammals are very close to the source.

Hearing Impairment and Other Physical Effects—It is unlikely that the sub-bottom profiler produces pulse levels strong enough to cause hearing impairment or other physical injuries even in an animal that is (briefly) in a position near the source. The sub-bottom profiler is usually operated simultaneously with other higher-power acoustic sources, including airguns. Many marine mammals will move away in response to the approaching higher-power sources or the vessel itself before the mammals would be close enough for there to be any possibility of effects from the less intense sounds from the sub-bottom profiler.

Heat-Flow Probe Deployment

During heat-flow measurements using a probe, the probe is a passive instrument and no noise is created by the mechanical action of the devices on the seafloor is not expected to be perceived by nearby fish and other marine organisms. Heat-flow measurement activities would be highly localized and short-term in duration and would not be expected to significantly interfere with marine mammal behavior. The potential direct effects include temporary localized disturbance or displacement from associated physical movement/actions of the operations. Additionally, the potential indirect effects may consist of very localized and transitory/short-term disturbance of bottom habitat and associated prey in shallow-water areas as a result of heat-flow probe measurements. NMFS believes that since the heat-flow probe is a passive instrument and has no mechanical action, it would not likely result in the harassment of marine mammals.

A maximum total of 200 heat-flow measurements would be obtained using these devices and ranging from 1 to 2 hours per measurement (for a total of approximately 320 hours of operations) and it is estimated that the pinger would operate continuously during each heat-flow probe deployment. The vessel would be stationary during heat-flow

probe deployment and repositioned to repeat the process, so the likelihood of a collision or entanglement with a marine mammal is very low. For the heat-flow measurements, the lance is 4.5 m and would disturb an area approximately 8 cm x 20 cm (3.1 in x 7.9 in). Assuming approximately 200 heat-flow measurements, the cumulative area of seafloor that could be disturbed during the proposed study would be approximately 32,000 cm² (4,960 in²).

Vessel Movement and Collisions

Vessel movement in the vicinity of marine mammals has the potential to result in either a behavioral response or a direct physical interaction. Both scenarios are discussed below in this section.

Behavioral Responses to Vessel Movement—There are limited data concerning marine mammal behavioral responses to vessel traffic and vessel noise, and a lack of consensus among scientists with respect to what these responses mean or whether they result in short-term or long-term adverse effects. In those cases where there is a busy shipping lane or where there is a large amount of vessel traffic, marine mammals (especially low frequency specialists) may experience acoustic masking (Hildebrand, 2005) if they are present in the area (*e.g.*, killer whales in Puget Sound; Foote *et al.*, 2004; Holt *et al.*, 2008). In cases where vessels actively approach marine mammals (*e.g.*, whale watching or dolphin watching boats), scientists have documented that animals exhibit altered behavior such as increased swimming speed, erratic movement, and active avoidance behavior (Bursk, 1983; Acevedo, 1991; Baker and MacGibbon, 1991; Trites and Bain, 2000; Williams *et al.*, 2002; Constantine *et al.*, 2003), reduced blow interval (Ritcher *et al.*, 2003), disruption of normal social behaviors (Lusseau, 2003, 2006), and the shift of behavioral activities which may increase energetic costs (Constantine *et al.*, 2003, 2004). A detailed review of marine mammal reactions to ships and boats is available in Richardson *et al.*, (1995). For each of the marine mammal taxonomy groups, Richardson *et al.*, (1995) provides the following assessment regarding reactions to vessel traffic:

Toothed whales—“In summary, toothed whales sometimes show no avoidance reaction to vessels, or even approach them. However, avoidance can occur, especially in response to vessels of types used to chase or hunt the animals. This may cause temporary displacement, but we know of no clear evidence that toothed whales have

abandoned significant parts of their range because of vessel traffic.”

Baleen whales—“When baleen whales receive low-level sounds from distant or stationary vessels, the sounds often seem to be ignored. Some whales approach the sources of these sounds. When vessels approach whales slowly and non-aggressively, whales often exhibit slow and inconspicuous avoidance maneuvers. In response to strong or rapidly changing vessel noise, baleen whales often interrupt their normal behavior and swim rapidly away. Avoidance is especially strong when a boat heads directly toward the whale.”

Behavioral responses to stimuli are complex and influenced by varying degrees by a number of factors, such as species, behavioral contexts, geographical regions, source characteristics (moving or stationary, speed, direction, etc.), prior experience of the animal and physical status of the animal. For example, studies have shown that beluga whales’ reaction varied when exposed to vessel noise and traffic. In some cases, beluga whales exhibited rapid swimming from ice-breaking vessels up to 80 km (43.2 nmi) away and showed changes in surfacing, breathing, diving, and group composition in the Canadian high Arctic where vessel traffic is rare (Finley *et al.*, 1990). In other cases, beluga whales were more tolerant of vessels, but responded differentially to certain vessels and operating characteristics by reducing their calling rates (especially older animals) in the St. Lawrence River where vessel traffic is common (Blane and Jaakson, 1994). In Bristol Bay, Alaska, beluga whales continued to feed when surrounded by fishing vessels and resisted dispersal even when purposefully harassed (Fish and Vania, 1971).

In reviewing more than 25 years of whale observation data, Watkins (1986) concluded that whale reactions to vessel traffic were “modified by their previous experience and current activity: Habituation often occurred rapidly, attention to other stimuli or preoccupation with other activities sometimes overcame their interest or wariness of stimuli.” Watkins noticed that over the years of exposure to ships in the Cape Cod area, minke whales changed from frequent positive interest (*e.g.*, approaching vessels) to generally uninterested reactions; fin whales changed from mostly negative (*e.g.*, avoidance) to uninterested reactions; fin whales changed from mostly negative (*e.g.*, avoidance) to uninterested reactions; right whales apparently continued the same variety of responses

(negative, uninterested, and positive responses) with little change; and humpbacks dramatically changed from mixed responses that were often negative to reactions that were often strongly positive. Watkins (1986) summarized that “whales near shore, even in regions with low vessel traffic, generally have become less wary of boats and their noises, and they have appeared to be less easily disturbed than previously. In particular locations with intense shipping and repeated approaches by boats (such as the whale-watching areas of Stellwagen Bank), more and more whales had positive reactions to familiar vessels, and they also occasionally approached other boats and yachts in the same ways.”

Although the radiated sound from the *Revelle* would be audible to marine mammals over a large distance, it is unlikely that marine mammals would respond behaviorally (in a manner that NMFS would consider harassment under the MMPA) to low-level distant shipping noise as the animals in the area are likely to be habituated to such noises (Nowacek *et al.*, 2004). In light of these facts, NMFS does not expect the *Revelle*’s movements to result in Level B harassment.

Vessel Strike—Ship strikes of cetaceans can cause major wounds, which may lead to the death of the animal. An animal at the surface could be struck directly by a vessel, a surfacing animal could hit the bottom of a vessel, or an animal just below the surface could be cut by a vessel’s propeller. The severity of injuries typically depends on the size and speed of the vessel (Knowlton and Kraus, 2001; Laist *et al.*, 2001; Vanderlaan and Taggart, 2007).

The most vulnerable marine mammals are those that spend extended periods of time at the surface in order to restore oxygen levels within their tissues after deep dives (*e.g.*, the sperm whale). In addition, some baleen whales, such as the North Atlantic right whale, seem generally unresponsive to vessel sound, making them more susceptible to vessel collisions (Nowacek *et al.*, 2004). These species are primarily large, slow moving whales. Smaller marine mammals (*e.g.*, bottlenose dolphins) move quickly through the water column and are often seen riding the bow wave of large ships. Marine mammal responses to vessels may include avoidance and changes in dive pattern (NRC, 2003).

An examination of all known ship strikes from all shipping sources (civilian and military) indicates vessel speed is a principal factor in whether a vessel strike results in death (Knowlton and Kraus, 2001; Laist *et al.*, 2001;

Jensen and Silber, 2003; Vanderlaan and Taggart, 2007). In assessing records in which vessel speed was known, Laist *et al.* (2001) found a direct relationship between the occurrence of a whale strike and the speed of the vessel involved in the collision. The authors concluded that most deaths occurred when a vessel was traveling in excess of 13 kts (24.1 km/hr, 14.9 mph).

SIO’s proposed operation of one source vessel for the proposed low-energy seismic survey is relatively small in scale (*i.e.*, a one vessel operation) compared to the number of other ships (*e.g.*, fishing, tourist, and other vessels) transiting at higher speeds in the same areas on an annual basis. The probability of vessel and marine mammal interactions occurring during the proposed low-energy seismic survey is unlikely due to the *Revelle*’s slow operational speed, which is typically 5 kts. Outside of seismic operations, the *Revelle*’s cruising speed would be approximately 10.1 to 14.5 kts, which is generally below the speed at which studies have noted reported increases of marine mammal injury or death (Laist *et al.*, 2001).

As a final point, the *Revelle* has a number of other advantages for avoiding ship strikes as compared to most commercial merchant vessels, including the following: The *Revelle*’s bridge and other observing stations offer good visibility to visually monitor for marine mammal presence; PSOs posted during operations scan the ocean for marine mammals and must report visual alerts of marine mammal presence to crew; and the PSOs receive extensive training that covers the fundamentals of visual observing for marine mammals and information about marine mammals and their identification at sea.

Entanglement

Entanglement can occur if wildlife becomes immobilized in survey lines, cables, nets, or other equipment that is moving through the water column. The proposed low-energy seismic survey would require towing approximately one 600 m cable streamers. While towing this size of an array carries some level of risk of entanglement for marine mammals due to the operational nature of the activity, entanglement is unlikely. Wildlife, especially slow moving individuals, such as large whales, have a low probability of becoming entangled due to slow speed of the survey vessel and onboard monitoring efforts. In May 2011, there was one recorded entrapment of an olive ridley sea turtle (*Lepidochelys olivacea*) in the R/V *Marcus G. Langseth*’s barovanes after the conclusion of a seismic survey off

Costa Rica. There have been cases of baleen whales, mostly gray whales (Heyning, 1990), becoming entangled in fishing lines. The probability for entanglement of marine mammals is considered very low because of the vessel speed and the monitoring efforts onboard the survey vessel. Furthermore, there has been no history of marine mammal entanglement with seismic equipment used by the U.S. academic research fleet.

The potential effects to marine mammals described in this section of the document do not take into consideration the proposed monitoring and mitigation measures described later in this document (see the "Proposed Mitigation" and "Proposed Monitoring and Reporting" sections) which, as noted are designed to effect the least practicable impact on affected marine mammal species and stocks.

Anticipated Effects on Marine Mammal Habitat

The proposed low-energy seismic survey is not anticipated to have any permanent impact on habitats used by the marine mammals in the proposed study area, including the food sources they use (*i.e.* fish and invertebrates). Additionally, no physical damage to any habitat is anticipated as a result of conducting airgun operations during the proposed low-energy seismic survey. While it is anticipated that the specified activity may result in marine mammals avoiding certain areas due to brief, temporary ensonification, this impact to habitat is temporary and was considered in further detail earlier in this document, as behavioral modification. The main impact associated with the proposed activity would be temporarily elevated noise levels and the associated direct effects on marine mammals in any particular area of the approximately 1,154 km² proposed study area, previously discussed in this notice.

The next section discusses the potential impacts of anthropogenic sound sources on common marine mammal prey in the proposed study area (*i.e.*, fish and invertebrates).

Anticipated Effects on Fish

One reason for the adoption of airguns as the standard energy source for marine seismic surveys is that, unlike explosives, they have not been associated with large-scale fish kills. However, existing information on the impacts of seismic surveys on marine fish and invertebrate populations is limited. There are three types of potential effects of exposure to seismic surveys: (1) Pathological, (2) physiological, and (3) behavioral.

Pathological effects involve lethal and temporary or permanent sub-lethal injury. Physiological effects involve temporary and permanent primary and secondary stress responses, such as changes in levels of enzymes and proteins. Behavioral effects refer to temporary and (if they occur) permanent changes in exhibited behavior (*e.g.*, startle and avoidance behavior). The three categories are interrelated in complex ways. For example, it is possible that certain physiological and behavioral changes could potentially lead to an ultimate pathological effect on individuals (*i.e.*, mortality).

The specific received sound levels at which permanent adverse effects to fish potentially could occur are little studied and largely unknown. Furthermore, the available information on the impacts of seismic surveys on marine fish is from studies of individuals or portions of a population; there have been no studies at the population scale. The studies of individual fish have often been on caged fish that were exposed to airgun pulses in situations not representative of an actual seismic survey. Thus, available information provides limited insight on possible real-world effects at the ocean or population scale. This makes drawing conclusions about impacts on fish problematic because, ultimately, the most important issues concern effects on marine fish populations, their viability, and their availability to fisheries.

Hastings and Popper (2005), Popper (2009), and Popper and Hastings (2009a,b) provided recent critical reviews of the known effects of sound on fish. The following sections provide a general synopsis of the available information on the effects of exposure to seismic and other anthropogenic sound as relevant to fish. The information comprises results from scientific studies of varying degrees of rigor plus some anecdotal information. Some of the data sources may have serious shortcomings in methods, analysis, interpretation, and reproducibility that must be considered when interpreting their results (see Hastings and Popper, 2005). Potential adverse effects of the program's sound sources on marine fish are noted.

Pathological Effects—The potential for pathological damage to hearing structures in fish depends on the energy level of the received sound and the physiology and hearing capability of the species in question. For a given sound to result in hearing loss, the sound must exceed, by some substantial amount, the hearing threshold of the fish for that sound (Popper, 2005). The consequences of temporary or permanent hearing loss in individual

fish on a fish population are unknown; however, they likely depend on the number of individuals affected and whether critical behaviors involving sound (*e.g.*, predator avoidance, prey capture, orientation and navigation, reproduction, etc.) are adversely affected.

Little is known about the mechanisms and characteristics of damage to fish that may be inflicted by exposure to seismic survey sounds. Few data have been presented in the peer-reviewed scientific literature. There are only two known papers with proper experimental methods, controls, and careful pathological investigation implicating sounds produced by actual seismic survey airguns in causing adverse anatomical effects. One such study indicated anatomical damage, and the second indicated TTS in fish hearing. The anatomical case is McCauley *et al.* (2003), who found that exposure to airgun sound caused observable anatomical damage to the auditory maculae of pink snapper (*Pagrus auratus*). This damage in the ears had not been repaired in fish sacrificed and examined almost two months after exposure. On the other hand, Popper *et al.* (2005) documented only TTS (as determined by auditory brainstem response) in two of three fish species from the Mackenzie River Delta. This study found that broad whitefish (*Coregonus nasus*) exposed to five airgun shots were not significantly different from those of controls. During both studies, the repetitive exposure to sound was greater than would have occurred during a typical seismic survey. However, the substantial low-frequency energy produced by the airguns (less than 400 Hz in the study by McCauley *et al.* [2003] and less than approximately 200 Hz in Popper *et al.* [2005]) likely did not propagate to the fish because the water in the study areas was very shallow (approximately nine m in the former case and less than two m in the latter). Water depth sets a lower limit on the lowest sound frequency that would propagate (the "cutoff frequency") at about one-quarter wavelength (Urick, 1983; Rogers and Cox, 1988).

Wardle *et al.* (2001) suggested that in water, acute injury and death of organisms exposed to seismic energy depends primarily on two features of the sound source: (1) The received peak pressure, and (2) the time required for the pressure to rise and decay. Generally, as received pressure increases, the period for the pressure to rise and decay decreases, and the chance of acute pathological effects increases. According to Buchanan *et al.*

(2004), for the types of seismic airguns and arrays involved with the proposed program, the pathological (mortality) zone for fish would be expected to be within a few meters of the seismic source. Numerous other studies provide examples of no fish mortality upon exposure to seismic sources (Falk and Lawrence, 1973; Holliday *et al.*, 1987; La Bella *et al.*, 1996; Santulli *et al.*, 1999; McCauley *et al.*, 2000a,b, 2003; Bjarti, 2002; Thomsen, 2002; Hassel *et al.*, 2003; Popper *et al.*, 2005; Boeger *et al.*, 2006).

An experiment of the effects of a single 700 in³ airgun was conducted in Lake Meade, Nevada (USGS, 1999). The data were used in an Environmental Assessment of the effects of a marine reflection survey of the Lake Meade fault system by the National Park Service (Paulson *et al.*, 1993, in USGS, 1999). The airgun was suspended 3.5 m (11.5 ft) above a school of threadfin shad in Lake Meade and was fired three successive times at a 30 second interval. Neither surface inspection nor diver observations of the water column and bottom found any dead fish.

For a proposed seismic survey in Southern California, USGS (1999) conducted a review of the literature on the effects of airguns on fish and fisheries. They reported a 1991 study of the Bay Area Fault system from the continental shelf to the Sacramento River, using a 10 airgun (5,828 in³) array. Brezzina and Associates were hired by USGS to monitor the effects of the surveys and concluded that airgun operations were not responsible for the death of any of the fish carcasses observed. They also concluded that the airgun profiling did not appear to alter the feeding behavior of sea lions, seals, or pelicans observed feeding during the seismic surveys.

Some studies have reported, some equivocally, that mortality of fish, fish eggs, or larvae can occur close to seismic sources (Kostyuchenko, 1973; Dalen and Knutsen, 1986; Booman *et al.*, 1996; Dalen *et al.*, 1996). Some of the reports claimed seismic effects from treatments quite different from actual seismic survey sounds or even reasonable surrogates. However, Payne *et al.* (2009) reported no statistical differences in mortality/morbidity between control and exposed groups of capelin eggs or monkfish larvae. Saetre and Ona (1996) applied a 'worst-case scenario' mathematical model to investigate the effects of seismic energy on fish eggs and larvae. They concluded that mortality rates caused by exposure to seismic surveys are so low, as compared to natural mortality rates, that the impact of seismic surveying on

recruitment to a fish stock must be regarded as insignificant.

Physiological Effects—Physiological effects refer to cellular and/or biochemical responses of fish to acoustic stress. Such stress potentially could affect fish populations by increasing mortality or reducing reproductive success. Primary and secondary stress responses of fish after exposure to seismic survey sound appear to be temporary in all studies done to date (Sverdrup *et al.*, 1994; Santulli *et al.*, 1999; McCauley *et al.*, 2000a,b). The periods necessary for the biochemical changes to return to normal are variable and depend on numerous aspects of the biology of the species and of the sound stimulus.

Behavioral Effects—Behavioral effects include changes in the distribution, migration, mating, and catchability of fish populations. Studies investigating the possible effects of sound (including seismic survey sound) on fish behavior have been conducted on both uncaged and caged individuals (*e.g.*, Chapman and Hawkins, 1969; Pearson *et al.*, 1992; Santulli *et al.*, 1999; Wardle *et al.*, 2001; Hassel *et al.*, 2003). Typically, in these studies fish exhibited a sharp startle response at the onset of a sound followed by habituation and a return to normal behavior after the sound ceased.

The former Minerals Management Service (MMS, 2005) assessed the effects of a proposed seismic survey in Cook Inlet. The seismic survey proposed using three vessels, each towing two four-airgun arrays ranging from 24,580.6 to 40,967.7 cm³ (1,500 to 2,500 in³). MMS noted that the impact to fish populations in the survey area and adjacent waters would likely be very low and temporary. MMS also concluded that seismic surveys may displace the pelagic fishes from the area temporarily when airguns are in use. However, fishes displaced and avoiding the airgun noise are likely to backfill the survey area in minutes to hours after cessation of seismic testing. Fishes not dispersing from the airgun noise (*e.g.*, demersal species) may startle and move short distances to avoid airgun emissions.

In general, any adverse effects on fish behavior or fisheries attributable to seismic testing may depend on the species in question and the nature of the fishery (season, duration, fishing method). They may also depend on the age of the fish, its motivational state, its size, and numerous other factors that are difficult, if not impossible, to quantify at this point, given such limited data on effects of airguns on fish, particularly under realistic at-sea conditions.

Anticipated Effects on Invertebrates

The existing body of information on the impacts of seismic survey sound on marine invertebrates is very limited. However, there is some unpublished and very limited evidence of the potential for adverse effects on invertebrates, thereby justifying further discussion and analysis of this issue. The three types of potential effects of exposure to seismic surveys on marine invertebrates are pathological, physiological, and behavioral. Based on the physical structure of their sensory organs, marine invertebrates appear to be specialized to respond to particle displacement components of an impinging sound field and not to the pressure component (Popper *et al.*, 2001).

The only information available on the impacts of seismic surveys on marine invertebrates involves studies of individuals; there have been no studies at the population scale. Thus, available information provides limited insight on possible real-world effects at the regional or ocean scale. The most important aspect of potential impacts concerns how exposure to seismic survey sound ultimately affects invertebrate populations and their viability, including availability to fisheries.

Literature reviews of the effects of seismic and other underwater sound on invertebrates were provided by Moriyasu *et al.* (2004) and Payne *et al.* (2008). The following sections provide a synopsis of available information on the effects of exposure to seismic survey sound on species of decapod crustaceans and cephalopods, the two taxonomic groups of invertebrates on which most such studies have been conducted. The available information is from studies with variable degrees of scientific soundness and from anecdotal information. A more detailed review of the literature on the effects of seismic survey sound on invertebrates is provided in Appendix D of NSF/USGS's PEIS (2011).

Pathological Effects—In water, lethal and sub-lethal injury to organisms exposed to seismic survey sound appears to depend on at least two features of the sound source: (1) The received peak pressure; and (2) the time required for the pressure to rise and decay. Generally, as received pressure increases, the period for the pressure to rise and decay decreases, and the chance of acute pathological effects increases. For the type of airgun array planned for the proposed program, the pathological (mortality) zone for crustaceans and cephalopods is

expected to be within a few meters of the seismic source, at most; however, very few specific data are available on levels of seismic signals that might damage these animals. This premise is based on the peak pressure and rise/decay time characteristics of seismic airgun arrays currently in use around the world.

Some studies have suggested that seismic survey sound has a limited pathological impact on early developmental stages of crustaceans (Pearson *et al.*, 1994; Christian *et al.*, 2003; DFO, 2004). However, the impacts appear to be either temporary or insignificant compared to what occurs under natural conditions. Controlled field experiments on adult crustaceans (Christian *et al.*, 2003, 2004; DFO, 2004) and adult cephalopods (McCauley *et al.*, 2000a,b) exposed to seismic survey sound have not resulted in any significant pathological impacts on the animals. It has been suggested that exposure to commercial seismic survey activities has injured giant squid (Guerra *et al.*, 2004), but the article provides little evidence to support this claim. Tenera Environmental (2011b) reported that Norris and Mohl (1983, summarized in Mariyasu *et al.*, 2004) observed lethal effects in squid (*Loligo vulgaris*) at levels of 246 to 252 dB after 3 to 11 minutes.

Andre *et al.* (2011) exposed four species of cephalopods (*Loligo vulgaris*, *Sepia officinalis*, *Octopus vulgaris*, and *Ilex coindetii*), primarily cuttlefish, to two hours of continuous 50 to 400 Hz sinusoidal wave sweeps at 157+/-5 dB re 1 μ Pa while captive in relatively small tanks. They reported morphological and ultrastructural evidence of massive acoustic trauma (*i.e.*, permanent and substantial alterations [lesions] of statocyst sensory hair cells) to the exposed animals that increased in severity with time, suggesting that cephalopods are particularly sensitive to low frequency sound. The received SPL was reported as 157+/-5 dB re 1 μ Pa, with peak levels at 175 dB re 1 μ Pa. As in the McCauley *et al.* (2003) paper on sensory hair cell damage in pink snapper as a result of exposure to seismic sound, the cephalopods were subjected to higher sound levels than they would be under natural conditions, and they were unable to swim away from the sound source.

Physiological Effects—Physiological effects refer mainly to biochemical responses by marine invertebrates to acoustic stress. Such stress potentially could affect invertebrate populations by increasing mortality or reducing reproductive success. Primary and

secondary stress responses (*i.e.*, changes in haemolymph levels of enzymes, proteins, etc.) of crustaceans have been noted several days or months after exposure to seismic survey sounds (Payne *et al.*, 2007). It was noted however, that no behavioral impacts were exhibited by crustaceans (Christian *et al.*, 2003, 2004; DFO, 2004). The periods necessary for these biochemical changes to return to normal are variable and depend on numerous aspects of the biology of the species and of the sound stimulus.

Behavioral Effects—There is increasing interest in assessing the possible direct and indirect effects of seismic and other sounds on invertebrate behavior, particularly in relation to the consequences for fisheries. Changes in behavior could potentially affect such aspects as reproductive success, distribution, susceptibility to predation, and catchability by fisheries. Studies investigating the possible behavioral effects of exposure to seismic survey sound on crustaceans and cephalopods have been conducted on both uncaged and caged animals. In some cases, invertebrates exhibited startle responses (*e.g.*, squid in McCauley *et al.*, 2000a,b). In other cases, no behavioral impacts were noted (*e.g.*, crustaceans in Christian *et al.*, 2003, 2004; DFO 2004). There have been anecdotal reports of reduced catch rates of shrimp shortly after exposure to seismic surveys; however, other studies have not observed any significant changes in shrimp catch rate (Andriquetto-Filho *et al.*, 2005). Similarly, Parry and Gason (2006) did not find any evidence that lobster catch rates were affected by seismic surveys. Any adverse effects on crustacean and cephalopod behavior or fisheries attributable to seismic survey sound depend on the species in question and the nature of the fishery (season, duration, fishing method). More information on the potential effects of airguns on fish and invertebrates are reviewed in section 3.2.4.3, section 3.3.4.3, and Appendix D of the NSF/USGS PEIS (2011).

Proposed Mitigation

In order to issue an Incidental Take Authorization (ITA) under section 101(a)(5)(D) of the MMPA, NMFS must set forth the permissible methods of taking pursuant to such activity, and other means of effecting the least practicable impact on such species or stock and its habitat, paying particular attention to rookeries, mating grounds, and areas of similar significance, and the availability of such species or stock

for taking for certain subsistence uses (where relevant).

SIO reviewed the following source documents and incorporated a suite of appropriate mitigation measures into the project description.

(1) Protocols used during previous NSF and USGS-funded seismic research cruises as approved by NMFS and detailed in the “Final Programmatic Environmental Impact Statement/ Overseas Environmental Impact Statement for Marine Seismic Research Funded by the National Science Foundation or Conducted by the U.S. Geological Survey;”

(2) Previous IHA applications and IHAs approved and authorized by NMFS; and

(3) Recommended best practices in Richardson *et al.* (1995), Pierson *et al.* (1998), and Weir and Dolman, (2007).

To reduce the potential for disturbance from acoustic stimuli associated with the activities, SIO proposed to implement the following mitigation measures for marine mammals:

- (1) Proposed exclusion zones around the sound source;
- (2) Speed and course alterations;
- (3) Shut-down procedures; and
- (4) Ramp-up procedures.

Proposed Exclusion Zones—During pre-planning of the cruise, the smallest airgun array was identified that could be used and still meet the geophysical scientific objectives. SIO use radii to designate exclusion and buffer zones and to estimate take for marine mammals. Table 2 (presented earlier in this document) shows the distances at which one would expect to receive three sound levels (160, 180, and 190 dB) from the two GI airgun array. The 180 and 190 dB level shut-down criteria are applicable to cetaceans and pinnipeds, respectively, as specified by NMFS (2000) and would be used to establish the exclusion and buffer zones.

Received sound levels have been modeled by L-DEO for a number of airgun configurations, including two 45 in³ Nucleus G airguns, in relation to distance and direction from the airguns (see Figure 2 of the IHA application). In addition, propagation measurements of pulses from two GI airguns have been reported for shallow water (approximately 30 m [98.4 ft] depth) in the Gulf of Mexico (Tolstoy *et al.*, 2004). However, measurements were not made for the two GI airguns in deep water. The model does not allow for bottom interactions, and is most directly applicable to deep water. Based on the modeling, estimates of the maximum distances from the GI airguns where sound levels are predicted to be 190,

180, and 160 dB re 1 μ Pa (rms) in intermediate and deep water were determined (see Table 2 above).

Empirical data concerning the 190, 180, and 160 dB (rms) distances were acquired for various airgun arrays based on measurements during the acoustic verification studies conducted by L-DEO in the northern Gulf of Mexico in 2003 (Tolstoy *et al.*, 2004) and 2007 to 2008 (Tolstoy *et al.*, 2009). Results of the 18 and 36 airgun arrays are not relevant for the two GI airguns to be used in the proposed low-energy seismic survey because the airgun arrays are not the same size or volume. The empirical data for the 6, 10, 12, and 20 airgun arrays indicate that, for deep water, the L-DEO model tends to overestimate the received sound levels at a given distance (Tolstoy *et al.*, 2004). Measurements were not made for the two GI airgun array in deep water; however, SIO proposes to use the safety radii predicted by L-DEO's model for the proposed GI airgun operations in intermediate and deep water, although they are likely conservative given the empirical results for the other arrays.

Based on the modeling data, the outputs from the pair of 45 in³ GI airguns proposed to be used during the low-energy seismic survey are considered a low-energy acoustic source in the NSF/USGS PEIS (2011) for marine seismic research. A low-energy seismic source was defined in the NSF/USGS PEIS as an acoustic source whose received level is less than or equal to 180 dB at 100 m (including any single or any two GI airguns and a single pair of clustered airguns with individual volumes of less than or equal to 250 in³). The NSF/USGS PEIS also established for these low-energy sources a standard exclusion zone of 100 m for all low-energy sources in water depths greater than 100 m. This standard 100 m exclusion zone would be used during the proposed low-energy seismic survey using the pair of 45 in³ GI airguns. The 180 and 190 dB (rms) radii are the current Level A harassment shut-down criteria applicable to cetaceans and pinnipeds, respectively; these levels were used to establish exclusion zones. Therefore, the assumed 180 and 190 dB radii are 100 m for intermediate and deep water. If the PSO detects a marine mammal within or about to enter the appropriate exclusion zone, the airguns would be shut down immediately.

Speed and Course Alterations—If a marine mammal is detected outside the exclusion zone and, based on its position and direction of travel (relative motion), is likely to enter the exclusion zone, changes of the vessel's speed and/or direct course would be considered if

this does not compromise operational safety or damage the deployed equipment. This would be done if operationally practicable while minimizing the effect on the planned science objectives. For marine seismic surveys towing large streamer arrays, course alterations are not typically implemented due to the vessel's limited maneuverability. However, the *Revelle* would be towing a relatively short hydrophone streamer, so its maneuverability during operations with the hydrophone streamer would not be limited as vessels towing long streamers, thus increasing the potential to implement course alterations, if necessary. After any such speed and/or course alteration is begun, the marine mammal activities and movements relative to the seismic vessel would be closely monitored to ensure that the marine mammal does not approach within the applicable exclusion zone. If the marine mammal appears likely to enter the exclusion zone, further mitigation actions would be taken, including further speed and/or course alterations, and/or shut-down of the airgun(s). Typically, during airgun operations, the source vessel is unable to change speed or course, and one or more alternative mitigation measures would need to be implemented.

Shut-Down Procedures—If a marine mammal is detected outside the exclusion zone for the airgun(s) but is likely to enter the exclusion zone, and the vessel's speed and/or course cannot be changed to avoid having the animal enter the exclusion zone, SIO would shut-down the operating airgun(s) before the animal is within the exclusion zone. Likewise, if a marine mammal is already within the exclusion zone when first detected, the airguns would be shut-down immediately.

Following a shut-down, SIO would not resume airgun activity until the marine mammal has cleared the exclusion zone, or until the PSO is confident that the animal has left the vicinity of the vessel. SIO would consider the animal to have cleared the exclusion zone if:

- A PSO has visually observed the animal leave the exclusion zone, or
- A PSO has not sighted the animal within the exclusion zone for 15 minutes for species with shorter dive durations (*i.e.*, small odontocetes and pinnipeds), or 30 minutes for species with longer dive durations (*i.e.*, mysticetes and large odontocetes, including sperm, dwarf and pygmy sperm, killer, and beaked whales).

Although power-down procedures are often standard operating practice for seismic surveys, they are not proposed

to be used during this planned low-energy seismic survey because powering down from two airguns to one airgun would make only a small difference in the exclusion zone(s) that probably would not be enough to allow continued one-airgun operations if a marine mammal came within the exclusion zone for two airguns.


Ramp-Up Procedures—Ramp-up of an airgun array provides a gradual increase in sound levels, and involves a step-wise increase in the number and total volume of airguns firing until the full volume of the airgun array is achieved. The purpose of a ramp-up is to “warn” marine mammals in the vicinity of the airguns and to provide the time for them to leave the area, avoiding any potential injury or impairment of their hearing abilities. SIO would follow a ramp-up procedure when the airgun array begins operating after a specified period without airgun operations or when a shut-down has exceeded that period. SIO proposes that, for the present cruise, this period would be approximately 15 minutes. SIO, L-DEO, USGS, NSF, and ASC have used similar periods (approximately 15 minutes) during previous low-energy seismic surveys.

Ramp-up would begin with a single GI airgun (45 in³). The second GI airgun (45 in³) would be added after 5 minutes. During ramp-up, the PSOs would monitor the exclusion zone, and if marine mammals are sighted, a shut-down would be implemented as though both GI airguns were operational.

If the complete exclusion zone has not been visible for at least 30 minutes prior to the start of operations in either daylight or nighttime, SIO would not commence the ramp-up. Given these provisions, it is likely that the airgun array would not be ramped-up from a complete shut-down during low light conditions, at night, or in thick fog, (*i.e.*, poor visibility conditions) because the outer part of the exclusion zone for that array would not be visible during those conditions. If one airgun has been operating, ramp-up to full power would be permissible during low light, at night, or in poor visibility, on the assumption that marine mammals would be alerted to the approaching seismic vessel by the sounds from the single airgun and could move away if they choose. SIO would not initiate a ramp-up of the airguns if a marine mammal is sighted within or near the applicable exclusion zones during day or night. NMFS refers the reader to Figure 2, which presents a flowchart representing the ramp-up and shut-down protocols described in this notice.

BILLING CODE 3510-22-P

Figure 2. Current mitigation procedures for low-energy seismic surveys.

**¹ Ramp-Up Procedures**

SIO has used similar periods (15 minutes) for previous low-energy seismic surveys. Ramp-up would not occur if a marine mammal has not cleared the exclusion zone for the full airgun array.

BILLING CODE 3510-22-C*Proposed Mitigation Conclusions*

NMFS has carefully evaluated the applicant's proposed mitigation measures and has considered a range of other measures in the context of ensuring that NMFS prescribes the means of effecting the least practicable impact on the affected marine mammal species and stocks and their habitat. NMFS's evaluation of potential measures included consideration of the

following factors in relation to one another:

- (1) The manner in which, and the degree to which, the successful implementation of the measure is expected to minimize adverse impacts to marine mammals;
- (2) The proven or likely efficacy of the specific measure to minimize adverse impacts as planned; and
- (3) The practicability of the measure for applicant implementation.

Any mitigation measure(s) prescribed by NMFS should be able to accomplish, have a reasonable likelihood of accomplishing (based on current science), or contribute to the accomplishment of one or more of the general goals listed below:

- (1) Avoidance of minimization of injury or death of marine mammals wherever possible (goals 2, 3, and 4 may contribute to this goal).
- (2) A reduction in the numbers of marine mammals (total number or

number at biologically important time or location) exposed to received levels of airguns, or other activities expected to result in the take of marine mammals (this goal may contribute to 1, above, or to reducing harassment takes only).

(3) A reduction in the number of time (total number or number at biologically important time or location) individuals would be exposed to received levels of airguns, or other activities expected to result in the take of marine mammals (this goal may contribute to 1, above, or to reducing harassment takes only).

(4) A reduction in the intensity of exposures (either total number or number at biologically important time or location) to received levels of airguns, or other activities, or other activities expected to result in the take of marine mammals (this goal may contribute to a, above, or to reducing the severity of harassment takes only).

(5) Avoidance or minimization of adverse effects to marine mammal habitat, paying special attention to the food base, activities that block or limit passage to or from biologically important areas, permanent destruction of habitat, or temporary destruction/disturbance of habitat during a biologically important time.

(6) For monitoring directly related to mitigation—an increase in the probability of detecting marine mammals, thus allowing for more effective implementation of the mitigation.

Based on NMFS's evaluation of the applicant's proposed measures, as well as other measures considered by NMFS or recommended by the public, NMFS has preliminarily determined that the proposed mitigation measures provide the means of effecting the least practicable impact on marine mammal species or stocks and their habitat, paying particular attention to rookeries, mating grounds, and areas of similar significance.

Proposed Monitoring and Reporting

In order to issue an ITA for an activity, section 101(a)(5)(D) of the MMPA states that NMFS must set forth "requirements pertaining to the monitoring and reporting of such taking." The MMPA implementing regulations at 50 CFR 216.104(a)(13) indicate that requests for IHAs must include the suggested means of accomplishing the necessary monitoring and reporting that would result in increased knowledge of the species and of the level of taking or impacts on populations of marine mammals that are expected to be present in the proposed action area. SIO submitted a marine mammal monitoring plan as part of the

IHA application. It can be found in Section 13 of the IHA application. The plan may be modified or supplemented based on comments or new information received from the public during the public comment period.

Monitoring measures prescribed by NMFS should accomplish one or more of the following general goals:

(1) An increase in the probability of detecting marine mammals, both within the mitigation zone (thus allowing for more effective implementation of the mitigation) and in general to generate more data to contribute to the analyses mentioned below;

(2) An increase in our understanding of how many marine mammals are likely to be exposed to levels of sound (airguns) that we associate with specific adverse effects, such as behavioral harassment, TTS, or PTS;

(3) An increase in our understanding of how marine mammals respond to stimuli expected to result in take and how anticipated adverse effects on individuals (in different ways and to varying degrees) may impact the population, species, or stock (specifically through effects on annual rates of recruitment or survival) through any of the following methods:

- Behavioral observations in the presence of stimuli compared to observations in the absence of stimuli (need to be able to accurately predict received level, distance from source, and other pertinent information);
- Physiological measurements in the presence of stimuli compared to observations in the absence of stimuli (need to be able to accurately predict received level, distance from source, and other pertinent information); and
- Distribution and/or abundance comparisons in times or areas with concentrated stimuli versus times or areas without stimuli;

(4) An increased knowledge of the affected species; and

(5) An increase in our understanding of the effectiveness of certain mitigation and monitoring measures.

Proposed Monitoring

SIO proposes to sponsor marine mammal monitoring during the proposed project, in order to implement the proposed mitigation measures that require real-time monitoring and to satisfy the anticipated monitoring requirements of the IHA. SIO's proposed "Monitoring Plan" is described below this section. The monitoring work described here has been planned as a self-contained project independent of any other related monitoring projects that may be occurring simultaneously in the same regions. SIO is prepared to

discuss coordination of their monitoring program with any related work that might be done by other groups insofar as this is practical and desirable.

Vessel-Based Visual Monitoring

PSOs would be based aboard the seismic source vessel and would watch for marine mammals near the vessel during daytime airgun operations and during any ramp-ups of the airguns at night. PSOs would also watch for marine mammals near the seismic vessel for at least 30 minutes prior to the start of airgun operations and after an extended shut-down (*i.e.*, greater than approximately 15 minutes for this proposed low-energy seismic survey). When feasible, PSOs would conduct observations during daytime periods when the seismic system is not operating (such as during transits) for comparison of sighting rates and behavior with and without airgun operations and between acquisition periods. Based on PSO observations, the airguns would be shut-down when marine mammals are observed within or about to enter a designated exclusion zone.

During airgun operations in the Southwest Pacific Ocean, East of New Zealand, at least three PSOs would be based aboard the *Revelle*. At least one PSO would stand watch at all times while the *Revelle* is operating airguns during the proposed low-energy seismic survey; this procedure would also be followed when the vessel is in transit. SIO would appoint the PSOs with NMFS's concurrence. The lead PSO would be experienced with marine mammal species in the Pacific Ocean and/or off the east coast of New Zealand, the second and third PSOs would receive additional specialized training from the lead PSO to ensure that they can identify marine mammal species commonly found in the Southwest Pacific Ocean. Observations would take place during ongoing daytime operations and ramp-ups of the airguns. During the majority of seismic operations, at least one PSO would be on duty from observation platforms (*i.e.*, the best available vantage point on the source vessel) to monitor marine mammals near the seismic vessel. PSO(s) would be on duty in shifts no longer than 4 hours in duration. Other crew would also be instructed to assist in detecting marine mammals and implementing mitigation requirements (if practical). Before the start of the low-energy seismic survey, the crew would be given additional instruction on how to do so.

The *Revelle* is a suitable platform for marine mammal observations and

would serve as the platform from which PSOs would watch for marine mammals before and during airgun operations. The *Revelle* has been used for marine mammal observations during the routine California Cooperative Oceanic Fisheries Investigations (CalCOFI). Two locations are likely as observation stations onboard the *Revelle*. Observing stations are located at the 02 level, with PSO eye level at approximately 10.4 m (34 ft) above the waterline and the PSO would have a good view around the entire vessel. At a forward-centered position on the 02 deck, the view is approximately 240° around the vessel; and one atop the aft hangar, with an aft-centered view includes the 100 m radius around the GI airguns. The PSO eye level on the bridge is approximately 15 m (49.2 ft) above sea level. PSOs would work on the enclosed bridge and adjoining aft steering station during any inclement weather.

Standard equipment for PSOs would be reticle binoculars and optical range finders. Night-vision equipment would be available at night and low-light conditions during the cruise. The PSOs would be in communication with ship's officers on the bridge and scientists in the vessel's operations laboratory, so they can advise promptly of the need for avoidance maneuvers or seismic source shut-down. During daylight, the PSO(s) would scan the area around the vessel systematically with reticle binoculars (e.g., 7 x 50 Fujinon FMTRC-SX), Big-eye binoculars (e.g., 25 x 150 Fujinon MT), optical range-finders (to assist with distance estimation), and the naked eye. These binoculars would have a built-in daylight compass. Estimating distances is done primarily with the reticles in the binoculars. The optical range-finders are useful in training PSOs to estimate distances visually, but are generally not useful in measuring distances to animals directly. At night, night-vision equipment would be available. The PSO(s) would be in direct (radio) wireless communication with ship's officers on the bridge and scientists in the vessel's operations laboratory during seismic operations, so they can advise the vessel operator, science support personnel, and the science party promptly of the need for avoidance maneuvers or a shut-down of the seismic source.

When a marine mammal is detected within or about to enter the designated exclusion zone, the airguns would immediately be shut-down, unless the vessel's speed and/or course can be changed to avoid having the animal enter the exclusion zone. The PSO(s) would continue to maintain watch to determine when the animal is outside

the exclusion zone by visual confirmation. Airgun operations would not resume until the animal is confirmed to have left the exclusion zone, or is not observed after 15 minutes for species with shorter dive durations (small odontocetes and pinnipeds) or 30 minutes for species with longer dive durations (mysticetes and large odontocetes, including sperm, dwarf and pygmy sperm, killer, and beaked whales).

PSO Data and Documentation

PSOs would record data to estimate the numbers of marine mammals exposed to various received sound levels and to document apparent disturbance reactions or lack thereof. Data would be used to estimate numbers of animals potentially "taken" by harassment. They would also provide information needed to order a shut-down of the airguns when a marine mammal is within or near the exclusion zone. Observations would also be made during daylight periods when the *Revelle* is underway without seismic airgun operations (i.e., transits to, from, and through the study area) to collect baseline biological data.

When a sighting is made, the following information about the sighting would be recorded:

1. Species, group size, age/size/sex categories (if determinable), behavior when first sighted and after initial sighting, heading (if consistent), bearing and distance from seismic vessel, sighting cue, apparent reaction to the seismic source or vessel (e.g., none, avoidance, approach, paralleling, etc.), and behavioral pace.
2. Time, location, heading, speed, activity of the vessel (including number of airguns operating and whether in state of ramp-up or shut-down), sea state, wind force, visibility, cloud cover, and sun glare.

The data listed under (2) would also be recorded at the start and end of each observation watch, and during a watch whenever there is a change in one or more of the variables.

All observations, as well as information regarding ramp-ups or shut-downs, would be recorded in a standardized format. Data would be entered into an electronic database. The data accuracy would be verified by computerized data validity checks as the data are entered and by subsequent manual checking of the database by the PSOs at sea. These procedures would allow initial summaries of data to be prepared during and shortly after the field program, and would facilitate transfer of the data to statistical,

graphical, and other programs for further processing and archiving.

Results from the vessel-based observations would provide the following information:

1. The basis for real-time mitigation (airgun shut-down).
2. Information needed to estimate the number of marine mammals potentially taken by harassment, which must be reported to NMFS.
3. Data on the occurrence, distribution, and activities of marine mammals in the area where the seismic study is conducted.
4. Information to compare the distance and distribution of marine mammals relative to the source vessel at times with and without airgun operations.
5. Data on the behavior and movement patterns of marine mammals seen at times with and without airgun operations.

Proposed Reporting

SIO would submit a comprehensive report to NMFS and NSF within 90 days after the end of the cruise. The report would describe the operations that were conducted and sightings of marine mammals near the operations. The report submitted to NMFS and NSF would provide full documentation of methods, results, and interpretation pertaining to all monitoring. The 90-day report would summarize the dates and locations of airgun operations and all marine mammal sightings (i.e., dates, times, locations, activities, and associated seismic survey activities). The report would include, at a minimum:

- Summaries of monitoring effort—total hours, total distances, and distribution of marine mammals through the study period accounting for Beaufort sea state and other factors affecting visibility and detectability of marine mammals;
- Analyses of the effects of various factors influencing detectability of marine mammals including Beaufort sea state, number of PSOs, and fog/glare;
- Species composition, occurrence, and distribution of marine mammal sightings including date, water depth, numbers, age/size/gender, and group sizes, and analyses of the effects of airgun operations;
- Sighting rates of marine mammals during periods with and without airgun operations (and other variables that could affect detectability);
- Initial sighting distances versus airgun operations state;
- Closest point of approach versus airgun operations state;

- Observed behaviors and types of movements versus airgun operations activity state;
- Numbers of sightings/individuals seen versus airgun operations state; and
- Distribution around the source vessel versus airgun operations state.

The report would also include estimates of the number and nature of exposures that could result in “takes” of marine mammals by harassment or in other ways. NMFS would review the draft report and provide any comments it may have, and SIO would incorporate NMFS’s comments and prepare a final report. After the report is considered final, it would be publicly available on the NMFS Web site at: <http://www.nmfs.noaa.gov/pr/permits/incidental/>.

Reporting Prohibited Take—In the unanticipated event that the specified activity clearly causes the take of a marine mammal in a manner prohibited by this IHA, such as an injury (Level A harassment), serious injury or mortality (e.g., ship-strike, gear interaction, and/or entanglement), SIO would immediately cease the specified activities and immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS at 301–427–8401 and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the following information:

- Time, date, and location (latitude/longitude) of the incident;
- Name and type of vessel involved;
- Vessel’s speed during and leading up to the incident;
- Description of the incident;
- Status of all sound source use in the 24 hours preceding the incident;
- Water depth;
- Environmental conditions (e.g., wind speed and direction, Beaufort sea state, cloud cover, and visibility);
- Description of all marine mammal observations in the 24 hours preceding the incident;
- Species identification or description of the animal(s) involved;
- Fate of the animal(s); and
- Photographs or video footage of the animal(s) (if equipment is available).

Activities shall not resume until NMFS is able to review the circumstances of the prohibited take. NMFS shall work with SIO to determine what is necessary to minimize the likelihood of further prohibited take and ensure MMPA compliance. SIO may not resume their activities until notified by NMFS via letter or email, or telephone.

Reporting an Injured or Dead Marine Mammal With an Unknown Cause of Death—In the event that SIO discover an injured or dead marine mammal, and

the lead PSO determines that the cause of the injury or death is unknown and the death is relatively recent (i.e., in less than a moderate state of decomposition), SIO shall immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301–427–8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the same information identified in the paragraph above. Activities may continue while NMFS reviews the circumstances of the incident. NMFS shall work with SIO to determine whether modifications in the activities are appropriate.

Reporting an Injured or Dead Marine Mammal Not Related to the Activities—In the event that SIO discovers an injured or dead marine mammal, and the lead PSO determines that the injury or death is not associated with or related to the activities authorized in the IHA (e.g., previously wounded animal, carcass with moderate or advanced decomposition, or scavenger damage), SIO shall report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301–427–8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov, within 24 hours of discovery. SIO shall provide photographs or video footage (if available) or other documentation of the stranded animal sighting to NMFS. Activities may continue while NMFS reviews the circumstances of the incident.

Estimated Take by Incidental Harassment

Except with respect to certain activities not pertinent here, the MMPA defines “harassment” as: Any act of pursuit, torment, or annoyance which (i) has the potential to injure a marine mammal or marine mammal stock in the wild [Level A harassment]; or (ii) has the potential to disturb a marine mammal or marine mammal stock in the wild by causing disruption of behavioral patterns, including, but not limited to, migration, breathing, nursing, breeding, feeding, or sheltering [Level B harassment].

TABLE 4—NMFS’S CURRENT UNDERWATER ACOUSTIC EXPOSURE CRITERIA

Criterion	Criterion definition	Threshold
Impulsive (non-explosive) sound		
Level A harassment (injury).	Permanent threshold shift (PTS) (Any level above that which is known to cause TTS).	180 dB re 1 μPa-m (root means square [rms]) (cetaceans) 190 dB re 1 μPa-m (rms) (pinnipeds)
Level B harassment.	Behavioral disruption (for impulsive noise).	160 dB re 1 μPa-m (rms)
Level B harassment.	Behavioral disruption (for continuous noise).	120 dB re 1 μPa-m (rms)

Level B harassment is anticipated and proposed to be authorized as a result of the proposed low-energy seismic survey in the Southwest Pacific Ocean, East of New Zealand. Acoustic stimuli (i.e., increased underwater sound) generated during the operation of the seismic airgun array are expected to result in the behavioral disturbance of some marine mammals. NMFS’s current underwater exposure criteria for impulsive sound are detailed in Table 4 (above). There is no evidence that the planned activities for which SIO seek the IHA could result in injury, serious injury, or mortality. The required mitigation and monitoring measures would minimize any potential risk for injury, serious injury, or mortality.

The following sections describe SIO’s methods to estimate take by incidental harassment and present the applicant’s estimates of the numbers of marine mammals that could be affected. The estimates are based on a consideration of the number of marine mammals that could be harassed during the approximately 135 hours and 1,250 km of seismic airgun operations with the two GI airgun array to be used.

There are no known systematic aircraft- or ship-based surveys conducted for marine mammals stock assessments and very limited population information available for marine mammals in offshore waters of the Southwest Pacific Ocean off the east coast of New Zealand. For most cetacean species, SIO and NMFS used densities from extensive NMFS Southwest Fisheries Science Center (SWFSC) cruises (Ferguson and Barlow, 2001, 2003; Barlow, 2003, 2010; Forney, 2007) in one province of Longhurst’s

(2006) pelagic biogeography, the California Current Province (CALC). That province is similar to the South Subtropical Convergence Province (SSTC) in which the proposed low-energy seismic survey is located, in that productivity is high and large pelagic fish such as tuna occur. Specifically, SIO and NMFS used the 1986 to 1996 data from blocks 35, 36, 47, 48, 59, and 60 of Ferguson and Barlow (2001, 2003), the 2001 data from Barlow (2003) for the Oregon, Washington, and California strata, and the 2005 and 2008 data from Forney (2007) and Barlow (2010), respectively, for the two strata combined. The densities used were effort-weighted means for the 10 locations (blocks or States). The surveys off California, Oregon, and Washington were conducted up to approximately 556 km (300.2 nmi) offshore, and most of those data were from offshore areas that overlap with the above blocks selected from Ferguson and Barlow (2001, 2003).

For pinnipeds, SIO and NMFS used the densities in Bonnell *et al.* (1992) of northern fur seals (*Callorhinus ursinus*) and northern elephant seals in offshore areas of the western U.S. (the only species regularly present in offshore areas there) to estimate the numbers of pinnipeds that might be present off New Zealand.

The marine mammal species that would be encountered during the

proposed low-energy seismic survey would be different from those sighted during surveys off the western U.S. and in the Eastern Tropical Pacific Ocean. However, the overall abundances of species groups with generally similar habitat requirements are expected to be roughly similar. Thus, SIO and NMFS used the data described above to estimate the group densities of beaked whales, delphinids, small whales, and mysticetes in the proposed study area. SIO and NMFS then estimated the relative abundance of individual southern species within the species groups using various surveys and other information from areas near the study area, and general information on species' distributions such as latitudinal ranges and group sizes. Group densities from northern species were multiplied by their estimated relative abundance off New Zealand divided by the relative abundance for all species in the species group to derive estimates for the southern species (see Table 3 of the IHA application).

Densities for several cetacean species are available for the Southern Ocean (Butterworth *et al.*, 1994), as follows: (1) For humpback, sei, fin, blue, sperm, killer, and pilot whales in Antarctic Management areas I to VI south of 60° South, based on the 1978/1979 to 1984 and 1985/1986 to 1990/1991 IWC/IDCR circumpolar sighting survey cruises, and

(2) for humpback, sei, fin, blue, and sperm whales extrapolated to latitudes 30 to 40° South, 40 to 50° South, 50 to 60° South based on Japanese scouting vessel data from 1965/1966 to 1977/1978 and 1978/1979 to 1987/1988. SIO and NMFS calculated densities based on abundance and surface areas given in Butterworth *et al.* (1994) and used the weighted or mean density for the Regions V and/or VI (whichever is available) due to locations that represent foraging areas or distributions for animals that are likely to move past New Zealand during northerly migrations or breed in New Zealand waters.

The densities used for purposes of estimating potential take do not take into account the patchy distributions of marine mammals in an ecosystem, at least on the moderate to fine scales over which they are known to occur. Instead, animals are considered evenly distributed throughout the assessed study area and seasonal movement patterns are not taken into account, as none are available. Although there is some uncertainty about the representativeness of the data and the assumptions used in the calculations below, the approach used here is believed to be the best available approach, using the best available science.

TABLE 5—ESTIMATED DENSITIES AND POSSIBLE NUMBER OF MARINE MAMMAL SPECIES THAT MIGHT BE EXPOSED TO GREATER THAN OR EQUAL TO 160 dB (AIRGUN OPERATIONS) DURING SIO'S PROPOSED LOW-ENERGY SEISMIC SURVEY (APPROXIMATELY 1,250 km OF TRACKLINES/APPROXIMATELY 1,154 km² ENSONIFIED AREA FOR AIRGUN OPERATIONS) IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND, MAY TO JUNE 2015

Species	Density U.S. West Coast/Southern Ocean/estimate used (number of animals/1,000 km ²) ¹	Calculated take from seismic airgun operations (<i>i.e.</i> , estimated number of individuals exposed to sound levels ≥160 dB re 1 μPa) ²	Proposed take authorization ³	Abundance ⁴	Approximate percentage of population estimate (proposed take) ⁵	Population trend ⁶
Mysticetes						
Southern right whale.	0.98/NA/0.98	1.13	2	8,000 to 15,000—Worldwide. 12,000—Southern Hemisphere. 2,700—Sub-Antarctic New Zealand.	0.03—Worldwide. 0.02—Southern Hemisphere. 0.07—Sub-Antarctic New Zealand.	Increasing at 7 to 8% per year.
Pygmy right whale.	0.39/NA/0.39	0.45	2	NA	NA	NA.
Humpback whale	0.98/0.25/0.25	0.29	2	35,000 to 42,000—Southern Hemisphere.	<0.01—Southern Hemisphere.	Increasing.
Antarctic minke whale.	0.59/NA/0.59	0.68	2	720,000 to 750,000—Southern Hemisphere.	<0.01—Southern Hemisphere.	Stable.

TABLE 5—ESTIMATED DENSITIES AND POSSIBLE NUMBER OF MARINE MAMMAL SPECIES THAT MIGHT BE EXPOSED TO GREATER THAN OR EQUAL TO 160 dB (AIRGUN OPERATIONS) DURING SIO'S PROPOSED LOW-ENERGY SEISMIC SURVEY (APPROXIMATELY 1,250 km OF TRACKLINES/APPROXIMATELY 1,154 km² ENSONIFIED AREA FOR AIRGUN OPERATIONS) IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND, MAY TO JUNE 2015—Continued

Species	Density U.S. West Coast/Southern Ocean/estimate used (number of animals/1,000 km ²) ¹	Calculated take from seismic airgun operations (i.e., estimated number of individuals exposed to sound levels ≥160 dB re 1 μPa) ²	Proposed take authorization ³	Abundance ⁴	Approximate percentage of population estimate (proposed take) ⁵	Population trend ⁶
Minke whale (including dwarf minke whale sub-species).	0.59/NA/0.59	0.68	2	720,000 to 750,000—Southern Hemisphere.	<0.01—Southern Hemisphere.	NA.
Bryde's whale	0.20/NA/0.20	0.23	2	At least 30,000 to 40,000—Worldwide. 21,000—Northwestern Pacific Ocean 48,109.	<0.01—Worldwide. <0.01—Northwestern Pacific Ocean <0.01.	NA.
Sei whale	0.59/0.08/0.08	0.09	2	80,000—Worldwide. 10,000—South of Antarctic Convergence.	<0.01—Worldwide. 0.02—South of Antarctic Convergence.	NA.
Fin whale	0.59/0.13/0.13	0.15	2	140,000—Worldwide. 15,000—South of Antarctic Convergence.	<0.01—Worldwide. 0.01—South of Antarctic Convergence.	NA.
Blue whale	0.59/0.05/0.05	0.06	2	8,000 to 9,000—Worldwide. 2,300—True Southern Hemisphere. 1,500—Pygmy.	0.03—Worldwide. 0.09—True Southern Hemisphere. 0.13—Pygmy.	NA.

Odontocetes

Sperm whale	1.62/1.16/1.16	1.34	10	360,000—Worldwide. 30,000—South of Antarctic Convergence.	<0.01—Worldwide. 0.03—South of Antarctic Convergence.	NA.
Pygmy sperm whale.	0.97/NA/0.97	1.12	5	NA	NA	NA.
Cuvier's beaked whale.	0.69/NA/0.69	0.80	2	600,000	<0.01	NA.
Shepherd's beaked whale.	0.46/NA/0.46	0.53	3	600,000	<0.01	NA.
Southern bottlenose whale.	0.46/NA/0.46	0.53	2	50,000—South of Antarctic Convergence 600,000.	<0.01—South of Antarctic Convergence <0.01.	NA.
Andrew's beaked whale.	0.46/NA/0.46	0.53	2	600,000	<0.01	NA.
Blainville's beaked whale.	0.23/NA/0.23	0.27	2	600,000	<0.01	NA.
Gray's beaked whale.	0.92/NA/0.92	1.06	2	600,000	<0.01	NA.
Hector's beaked whale.	0.46/NA/0.46	0.53	2	600,000	<0.01	NA.
Spade-toothed beaked whale.	0.23/NA/0.23	0.27	2	600,000	<0.01	NA.
Strap-toothed beaked whale.	0.69/NA/0.69	0.80	3	600,000	<0.01	NA.
Killer whale	0.45/5.70/5.70	6.58	12	80,000—South of Antarctic Convergence.	0.02—South of Antarctic Convergence.	NA.
False killer whale	0.27/NA/0.27	0.31	10	NA	NA	NA.

TABLE 5—ESTIMATED DENSITIES AND POSSIBLE NUMBER OF MARINE MAMMAL SPECIES THAT MIGHT BE EXPOSED TO GREATER THAN OR EQUAL TO 160 dB (AIRGUN OPERATIONS) DURING SIO'S PROPOSED LOW-ENERGY SEISMIC SURVEY (APPROXIMATELY 1,250 km OF TRACKLINES/APPROXIMATELY 1,154 km² ENSONIFIED AREA FOR AIRGUN OPERATIONS) IN THE SOUTHWEST PACIFIC OCEAN, EAST OF NEW ZEALAND, MAY TO JUNE 2015—Continued

Species	Density U.S. West Coast/Southern Ocean/estimate used (number of animals/1,000 km ²) ¹	Calculated take from seismic airgun operations (<i>i.e.</i> , estimated number of individuals exposed to sound levels ≥160 dB re 1 μPa) ²	Proposed take authorization ³	Abundance ⁴	Approximate percentage of population estimate (proposed take) ⁵	Population trend ⁶
Long-finned pilot whale.	0.27/6.41/6.41	7.40	20	200,000—South of Antarctic Convergence.	0.01—South of Antarctic Convergence.	NA.
Short-finned pilot whale.	0.45/NA/0.45	0.52	20	At least 600,000—Worldwide	<0.01—Worldwide.	NA.
Bottlenose dolphin.	81.55/NA/81.55	94.11	95	At least 614,000—Worldwide	0.02—Worldwide	NA.
Dusky dolphin	81.55/NA/81.55	94.11	95	12,000 to 20,000—New Zealand	0.79—New Zealand.	NA.
Hector's dolphin	32.62/NA/32.62	37.64	38	7,400	0.51	Declining.
Hourglass dolphin.	48.93/NA/48.93	56.47	57	144,000 to 150,000—South of Antarctic Convergence.	0.04—South of Antarctic Convergence.	NA.
Short-beaked common dolphin.	163.10/NA/163.10	188.22	189	At least 3,500,000—Worldwide	<0.01—Worldwide.	NA.
Southern right whale dolphin.	48.93/NA/48.93	56.46	57	NA	NA	NA.
Pinnipeds						
Southern elephant seal.	5.11/NA/5.11	5.90	6	640,000 to 650,000—Worldwide. 470,000—South Georgia Island 607,000.	<0.01—Worldwide or South Georgia Island.	Increasing, decreasing, or stable depending on breeding population.
New Zealand fur seal.	12.79/NA/12.79	14.76	15	135,000—Worldwide. 50,000 to 100,000—New Zealand.	0.01—Worldwide. 0.03—New Zealand.	Increasing.

NA = Not available or not assessed.

¹ Densities based on sightings from NMFS SWFSC, IWC, and Bonnell *et al.* (2012) data.

² Calculated take is estimated density multiplied by the area ensonified to 160 dB (rms) around the proposed seismic tracklines, increased by 25% for contingency.

³ Adjusted to account for average group size.

⁴ See population estimates for marine mammal species in Table 3 (above).

⁵ Total proposed authorized takes expressed as percentages of the species or regional populations.

⁶ Jefferson *et al.* (2008).

Numbers of marine mammals that might be present and potentially disturbed are estimated based on the available data about marine mammal distribution and densities in the U.S. west coast and Southern Ocean as a proxy for the proposed study area off the east coast of New Zealand. SIO estimated the number of different individuals that may be exposed to airgun sounds with received levels greater than or equal to 160 dB re 1 μPa (rms) for seismic airgun operations on one or more occasions by considering the total marine area that would be within the 160 dB radius around the operating airgun array on at least one

occasion and the expected density of marine mammals in the area (in the absence of the low-energy seismic survey). The number of possible exposures can be estimated by considering the total marine area that would be within the 160 dB radius (the diameter is 400 m multiplied by 2 for deep water depths, the diameter is 600 m multiplied by 2 for intermediate water depths) around the operating airguns, including areas of overlap. The spacing of tracklines is 500 m (1,640.4 ft) in the smaller grids and 1,250 m (4,101.1 ft) in the larger grids. Overlap was measured using GIS and was minimal (area with overlap is equal to

1.13 multiplied by the area without overlap). The take estimates were calculated without overlap. The 160 dB radii are based on acoustic modeling data for the airguns that may be used during the proposed action (see SIO's IHA application). During the proposed low-energy seismic survey, the transect lines are widely spaced relative to the 160 dB distance. As summarized in Table 2 (see Table 1 and Figure 2 of the IHA application), the modeling results for the proposed low-energy seismic airgun array indicate the received levels are dependent on water depth. Since the majority of the proposed airgun operations would be conducted in

waters 100 to 1,000 m deep or greater than 1,000 m deep, the buffer zone of 600 m or 400 m, respectively, for the two 45 in³ GI airguns was used.

The number of different individuals potentially exposed to received levels greater than or equal to 160 dB re 1 μ Pa (rms) from seismic airgun operations was calculated by multiplying:

(1) The expected species density (in number/km²), times.

(2) The anticipated area to be ensonified to that level during airgun operations (excluding overlap).

The area expected to be ensonified was determined by entering the planned tracklines into MapInfo GIS using the GIS to identify the relevant areas by "drawing" the applicable 160 dB (rms) isopleth around each trackline, and then calculating the total area within the isopleth. Applying the approach described above, approximately 1,153.6 km² (including the 25% contingency [approximately 923 km² without contingency]) would be ensonified within the 160 dB isopleth for seismic airgun operations on one or more occasions during the proposed low-energy seismic survey. The total ensonified area (1,154 km² [336.5 nmi²]) was calculated by adding 847 km² (246.9 nmi²) in deep water, 76 km² (22.2 nmi²), and 230.8 km² (67.3 nmi²) for the 25% contingency. The take calculations within the study sites do not explicitly add animals to account for the fact that new animals (*i.e.*, turnover) not accounted for in the initial density snapshot could also approach and enter the area ensonified above 160 dB for seismic airgun operations. However, studies suggest that many marine mammals would avoid exposing themselves to sounds at this level, which suggests that there would not necessarily be a large number of new animals entering the area once the seismic survey started. Because this approach for calculating take estimates does not account for turnover in the marine mammal populations in the area during the course of the proposed low-energy seismic survey, the actual number of individuals exposed may be underestimated. However, any underestimation is likely offset by the conservative (*i.e.*, probably overestimated) line-kilometer distances (including the 25% contingency) used to calculate the survey area, and the fact the approach assumes that no cetaceans or pinnipeds would move away or toward the tracklines as the *Revelle* approaches in response to increasing sound levels before the levels reach 160 dB for seismic airgun operations, which is likely to occur and which would decrease the density of marine

mammals in the survey area. Another way of interpreting the estimates in Table 6 is that they represent the number of individuals that would be expected (in absence of a seismic program) to occur in the waters that would be exposed to greater than or equal to 160 dB (rms) for seismic airgun operations.

SIO's estimates of exposures to various sound levels assume that the proposed low-energy seismic survey would be carried out in full; however, the ensonified areas calculated using the planned number of line-kilometers has been increased by 25% to accommodate lines that may need to be repeated, equipment testing, etc. As is typical during offshore seismic surveys, inclement weather and equipment malfunctions would be likely to cause delays and may limit the number of useful line-kilometers of airgun operations that can be undertaken. The estimates of the numbers of marine mammals potentially exposed to 160 dB (rms) received levels are precautionary and probably overestimate the actual numbers of marine mammals that could be involved. These estimates assume that there would be no weather, equipment, or mitigation delays that limit the airgun operations, which is highly unlikely.

Table 5 shows the estimates of the number of different individual marine mammals anticipated to be exposed to greater than or equal to 160 dB re 1 μ Pa (rms) for seismic airgun operations during the low-energy seismic survey if no animals moved away from the survey vessel. The total proposed take authorization is given in the column that is fourth from the left of Table 5.

Encouraging and Coordinating Research

SIO and NSF would coordinate the planned marine mammal monitoring program associated with the proposed low-energy seismic survey with other parties that express interest in this activity and area. SIO and NSF would coordinate with applicable U.S. agencies (*e.g.*, NMFS) and the government of New Zealand, and would comply with their requirements. The proposed low-energy seismic survey falls under Level 3 of the "Code of Conduct for minimizing acoustic disturbance to marine mammals from seismic survey operations" issued by New Zealand. Level 3 seismic surveys are exempt from the provisions of the Code of Conduct.

Impact on Availability of Affected Species or Stock for Taking for Subsistence Uses

Section 101(a)(5)(D) of the MMPA also requires NMFS to determine that the authorization would not have an unmitigable adverse impact on the availability of marine mammal species or stocks for subsistence use. There are no relevant subsistence uses of marine mammals implicated by this action (in the Southwest Pacific Ocean, East of New Zealand study area). Therefore, NMFS has determined that the total taking of affected species or stocks would not have an unmitigable adverse impact on the availability of such species or stocks for taking for subsistence purposes.

Analysis and Preliminary Determinations

Negligible Impact

Negligible impact is "an impact resulting from the specified activity that cannot be reasonably expected to, and is not reasonably likely to, adversely affect the species or stock through effects on annual rates of recruitment or survival" (50 CFR 216.103). A negligible impact finding is based on the lack of likely adverse effects on annual rates of recruitment or survival (*i.e.*, population-level effects). An estimate of the number of Level B harassment takes, alone, is not enough information on which to base an impact determination. In addition to considering estimates of the number of marine mammals that might be "taken" through behavioral harassment, NMFS must consider other factors, such as the likely nature of any responses (their intensity, duration, etc.) and the context of any responses (critical reproductive time or location, migration, etc.), as well as the number and nature of estimated Level A harassment takes, the number of estimated mortalities, effects on habitat, and the status of the species.

In making a negligible impact determination, NMFS evaluated factors such as:

- (1) The number of anticipated serious injuries and or mortalities;
- (2) The number and nature of anticipated injuries;
- (3) The number, nature, intensity, and duration of takes by Level B harassment (all of which are relatively limited in this case);
- (4) The context in which the takes occur (*e.g.*, impacts to areas of significance, impacts to local populations, and cumulative impacts when taking into account successive/contemporaneous actions when added to baseline data);

(5) The status of stock or species of marine mammals (*i.e.*, depleted, not depleted, decreasing, increasing, stable, impact relative to the size of the population);

(6) Impacts on habitat affecting rates of recruitment/survival; and

(7) The effectiveness of monitoring and mitigation measures.

NMFS has preliminarily determined that the specified activities associated with the marine seismic survey are not likely to cause PTS, or other (non-auditory) injury, serious injury, or death, based on the analysis above and the following factors:

(1) The likelihood that, given sufficient notice through relatively slow ship speed, marine mammals are expected to move away from a noise source that is annoying prior to its becoming potentially injurious;

(2) The availability of alternate areas of similar habitat value for marine mammals to temporarily vacate the survey area during the operation of the airgun(s) to avoid acoustic harassment;

(3) The potential for temporary or permanent hearing impairment is relatively low and would likely be avoided through the implementation of the required monitoring and mitigation measures (including shut-down measures); and

(4) The likelihood that marine mammal detection ability by trained PSOs is high at close proximity to the vessel.

No injuries, serious injuries, or mortalities are anticipated to occur as a result of the SIO's planned low-energy seismic survey, and none are proposed to be authorized by NMFS. Table 5 of this document outlines the number of requested Level B harassment takes that are anticipated as a result of these activities. Due to the nature, degree, and context of Level B (behavioral) harassment anticipated and described in this notice (see "Potential Effects on Marine Mammals" section above), the activity is not expected to impact rates of annual recruitment or survival for any affected species or stock, particularly given NMFS's and the applicant's proposed mitigation, monitoring, and reporting measures to minimize impacts to marine mammals. Additionally, the low-energy seismic survey would not adversely impact marine mammal habitat.

Many animals perform vital functions, such as feeding, resting, traveling, and socializing, on a diel cycle (*i.e.*, 24 hr cycle). Behavioral reactions to noise exposure (such as disruption of critical life functions, displacement, or avoidance of important habitat) are more likely to be significant if they last

more than one diel cycle or recur on subsequent days (Southall *et al.*, 2007). While airgun operations are anticipated to occur on consecutive days, the estimated duration of the survey would not last more than a total of approximately 27 operational days. Additionally, the low-energy seismic survey would be increasing sound levels in the marine environment in a relatively small area surrounding the vessel (compared to the range of the animals), which is constantly travelling over distances, so individual animals likely would only be exposed to and harassed by sound for less than a day.

As mentioned previously, NMFS estimates that 32 species of marine mammals under its jurisdiction could be potentially affected by Level B harassment over the course of the IHA. The population estimates for the marine mammal species that may be taken by Level B harassment were provided in Table 3 and 5 of this document. As shown in those tables, the proposed takes represent small proportions of the overall populations of these marine mammal species where abundance estimates are available (*i.e.*, less than 1%).

Of the 32 marine mammal species under NMFS jurisdiction that may or are known to likely occur in the study area, six are listed as threatened or endangered under the ESA: Southern right, humpback, sei, fin, blue, and sperm whales. These species are also considered depleted under the MMPA. None of the other marine mammal species that may be taken are listed as depleted under the MMPA. Of the ESA-listed species, incidental take has been requested to be authorized for six species. As mitigation to reduce impacts to the affected species or stocks, SIO would be required to cease airgun operations if any marine mammal enters designated exclusion zones. No injury, serious injury, or mortality is expected to occur for any of these species, and due to the nature, degree, and context of the Level B harassment anticipated, and the activity is not expected to impact rates of recruitment or survival for any of these species.

NMFS has preliminarily determined that, provided that the aforementioned mitigation and monitoring measures are implemented, the impact of conducting a low-energy marine seismic survey in the Southwest Pacific Ocean, May to June 2015, may result, at worst, in a modification in behavior and/or low-level physiological effects (Level B harassment) of certain species of marine mammals.

While behavioral modifications, including temporarily vacating the area

during the operation of the airgun(s), may be made by these species to avoid the resultant acoustic disturbance, the availability of alternate areas for species to move to and the short and sporadic duration of the research activities, have led NMFS to preliminarily determine that the taking by Level B harassment from the specified activity would have a negligible impact on the affected species in the specified geographic region. Due to the nature, degree, and context of Level B (behavioral) harassment anticipated and described (see "Potential Effects on Marine Mammals" section above) in this notice, the proposed activity is not expected to impact rates of annual recruitment or survival for any affected species or stock, particularly given the NMFS and applicant's proposal to implement mitigation and monitoring measures would minimize impacts to marine mammals. Based on the analysis contained herein of the likely effects of the specified activity on marine mammals and their habitat, and taking into consideration the implementation of the proposed monitoring and mitigation measures, NMFS preliminarily finds that the total marine mammal take from SIO's proposed low-energy seismic survey would have a negligible impact on the affected marine mammal species or stocks.

Small Numbers

As mentioned previously, NMFS estimates that 32 species of marine mammals under its jurisdiction could be potentially affected by Level B harassment over the course of the IHA. The population estimates for the marine mammal species that may be taken by Level B harassment were provided in Tables 3 and 5 of this document.

The estimated numbers of individual cetaceans and pinnipeds that could be exposed to seismic sounds with received levels greater than or equal to 160 dB re 1 μ Pa (rms) during the proposed low-energy seismic survey (including a 25% contingency) are in Table 5 of this document. Of the cetaceans, 2 southern right, 2 pygmy right, 2 humpback, 2 Antarctic minke, 2 minke, 2 Bryde's, 2 sei, 2 fin, 2 blue, and 10 sperm whales could be taken by Level B harassment during the proposed low-energy seismic survey, which would represent 0.03, unknown, 0.1, less than 0.01, 0.03, and 0.03% of the affected worldwide or regional populations, respectively. In addition, 5 pygmy sperm, 2 Cuvier's beaked, 3 Shepherd's beaked, 2 southern bottlenose, 2 Andrew's beaked, 2 Blainville's beaked, 2 Gray's beaked,

m) water depths. Airgun operations would take approximately 135 hours in total and 1,250 km, and the remainder of the time would be spent in transit and collecting heat-flow measurements and sediment core samples. The low-energy seismic survey would be conducted as specified in SIO's IHA application and the associated NSF and SIO Environmental Analysis.

3. This Authorization does not permit incidental takes of marine mammals in the territorial sea of foreign nations, as the MMPA does not apply in those waters. The territorial sea extends at the most 22.2 kilometers (km) (12 nautical miles [nmi]) from the baseline of a coastal State.

4. Species Authorized and Level of Takes

(a) The incidental taking of marine mammals, by Level B harassment only, is limited to the following species in the waters of the Southwest Pacific Ocean, East of New Zealand:

(i) *Mysticetes*—see Table 5 (above) for authorized species and take numbers.

(ii) *Odontocetes*—see Table 5 (above) for authorized species and take numbers.

(iii) *Pinnipeds*—see Table 5 (above) for authorized species and take numbers.

(iv) If any marine mammal species are encountered during seismic activities that are not listed in Table 5 (above) for authorized taking and are likely to be exposed to sound pressure levels (SPLs) greater than or equal to 160 dB re 1 μ Pa (rms) for seismic airgun operations, then the SIO must alter speed or course or shut-down the airguns to prevent take.

(b) The taking by injury (Level A harassment), serious injury, or death of any of the species listed in Condition 4(a) above or the taking of any kind of any other species of marine mammal is prohibited and may result in the modification, suspension, or revocation of this Authorization.

5. The sources authorized for taking by Level B harassment are limited to the following acoustic sources, absent an amendment to this Authorization:

A two Generator Injector (GI) airgun array (each with a discharge volume of 45 cubic inches [in³]) with a total volume of 90 in³ (or smaller).

6. Prohibited Take

The taking of any marine mammal in a manner prohibited under this Authorization must be reported immediately to the Office of Protected Resources, National Marine Fisheries Service (NMFS), at 301-427-8401.

7. Mitigation and Monitoring Requirements

The SIO is required to implement the following mitigation and related

monitoring requirements when conducting the specified activities to achieve the least practicable impact on affected marine mammal species or stocks:

Protected Species Observers and Visual Monitoring

(a) Utilize at least one NMFS-qualified, vessel-based Protected Species Observer (PSO) to visually watch for and monitor marine mammals near the seismic source vessel during daylight airgun operations (from nautical twilight-dawn to nautical twilight-dusk) and before and during ramp-ups of airguns day or night. Three PSOs shall be based onboard the vessel.

(i) The *Revelle's* vessel crew shall also assist in detecting marine mammals, when practicable.

(ii) PSOs shall have access to reticle binoculars (7 x 50 Fujinon) equipped with a built-in daylight compass and range reticles, big-eye binoculars (25 x 150), optical range finders, and night-vision devices.

(iii) PSO shifts shall last no longer than 4 hours at a time.

(iv) PSO(s) shall also make observations during daylight periods when the seismic airguns are not operating, when feasible, for comparison of animal abundance and behavior.

(v) PSO(s) shall conduct monitoring while the airgun array and streamer(s) are being deployed or recovered from the water.

(b) PSO(s) shall record the following information when a marine mammal is sighted:

(i) Species, group size, age/size/sex categories (if determinable), behavior when first sighted and after initial sighting, heading (if consistent), bearing and distance from seismic vessel, sighting cue, apparent reaction to the airguns or vessel (e.g., none, avoidance, approach, paralleling, etc., and including responses to ramp-up), and behavioral pace; and

(ii) Time, location, heading, speed, activity of the vessel (including number of airguns operating and whether in state of ramp-up or shut-down), Beaufort sea state and wind force, visibility, and sun glare; and

(iii) The data listed under Condition 7(b)(ii) shall also be recorded at the start and end of each observation watch and during a watch whenever there is a change in one or more of the variables.

Buffer and Exclusion Zones

(c) Establish a 160 dB re 1 μ Pa (rms) buffer zone, as well as a 180 dB re 1 μ Pa (rms) exclusion zone for cetaceans and a 190 dB re 1 μ Pa (rms) exclusion zone

for pinnipeds before the two GI airgun array (90 in³ total volume) is in operation. See Table 2 (above) for distances and buffer and exclusion zones.

Visual Monitoring at the Start of the Airgun Operations

(d) Visually observe the entire extent of the exclusion zone (180 dB re 1 μ Pa [rms] for cetaceans and 190 dB re 1 μ Pa [rms] for pinnipeds; see Table 2 [above] for distances) using two NMFS-qualified PSOs, for at least 30 minutes prior to starting the airgun array (day or night).

(i) If the PSO(s) sees a marine mammal within the exclusion zone, SIO must delay the seismic survey until the marine mammal(s) has left the area. If the PSO(s) sees a marine mammal that surfaces, then dives below the surface, the PSO(s) shall continue to observe the exclusion zone for 30 minutes, and if the PSO sees no marine mammals during that time, the PSO should assume that the animal has moved beyond the exclusion zone.

(ii) If for any reason the entire radius cannot be seen for the entire 30 minutes (i.e., rough seas, fog, darkness), or if marine mammals are near, approaching, or in the exclusion zone, the airguns may not be ramped-up. If one airgun is already running at a source level of at least 180 dB re 1 μ Pa (rms), SIO may start the second airgun without observing the entire exclusion zone for 30 minutes prior, provided no marine mammals are known to be near the exclusion zone (in accordance with Condition 7[e] below).

Ramp-Up Procedures

(e) Implement a "ramp-up" procedure, which means starting with a single GI airgun and adding a second GI airgun after five minutes, when starting up at the beginning of seismic operations or anytime after the entire array has been shut-down for more than 15 minutes. During ramp-up, the two PSOs shall monitor the exclusion zone, and if marine mammals are sighted, a shut-down shall be implemented as though the full array (both GI airguns) were operational. Therefore, initiation of ramp-up procedures from shut-down requires that the two PSOs be able to view the full exclusion zone as described in Condition 7(d) (above).

Shut-Down Procedures

(f) Shut-down the airgun(s) if a marine mammal is detected within, approaches, or enters the relevant exclusion zone (as defined in Table 2, above). A shut-down means all operating airguns are shut-down (i.e., turned off).

(g) Following a shut-down, the airgun activity shall not resume until the PSO(s) has visually observed the marine mammal(s) exiting the exclusion zone and determined it is not likely to return, or has not seen the marine mammal within the exclusion zone for 15 minutes, for species with shorter dive durations (small odontocetes and pinnipeds), or 30 minutes for species with longer dive durations (mysticetes and large odontocetes, including sperm, dwarf and pygmy sperm, killer, and beaked whales).

(h) Following a shut-down and subsequent animal departure, airgun operations may resume, following the ramp-up procedures described in Condition 7(e).

Speed or Course Alteration

(i) Alter speed or course during seismic operations if a marine mammal, based on its position and relative motion, appears likely to enter the relevant exclusion zone. If speed or course alteration is not safe or practicable, or if after alteration the marine mammal still appears likely to enter the exclusion zone, further mitigation measures, such as a shut-down, shall be taken.

Survey Operations During Low-Light Hours

(j) Marine seismic surveying may continue into low-light hours if such segment(s) of the survey is initiated when the entire relevant exclusion zones are visible and can be effectively monitored.

(k) No initiation of airgun array operations is permitted from a shut-down position during low-light hours (such as in dense fog or heavy rain) when the entire relevant exclusion zone cannot be effectively monitored by the PSO(s) on duty.

(l) To the maximum extent practicable, schedule seismic operations (*i.e.*, shooting airguns) during daylight hours, and heat-flow measurements at nighttime hours.

8. Reporting Requirements

SIO are required to:

(a) Submit a draft report on all activities and monitoring results to the Office of Protected Resources, NMFS, within 90 days of the completion of the *Revelle's* Southwest Pacific Ocean, East of New Zealand cruise. This report must contain and summarize the following information:

(i) Dates, times, locations, heading, speed, weather, sea conditions (including Beaufort sea state and wind force), and associated activities during all seismic operations and marine mammal sightings;

(ii) Species, number, location, distance from the vessel, and behavior of any marine mammals, as well as associated seismic activity (*e.g.*, number of shut-downs), observed throughout all monitoring activities.

(iii) An estimate of the number (by species) of marine mammals that: (A) Are known to have been exposed to the seismic activity (based on visual observation) at received levels greater than or equal to 160 dB re 1 μ Pa (rms) (for seismic airgun operations), and/or 180 dB re 1 μ Pa (rms) for cetaceans and 190 dB re 1 μ Pa (rms) for pinnipeds, with a discussion of any specific behaviors those individuals exhibited; and (B) may have been exposed (based on modeled values for the two GI airgun array) to the seismic activity at received levels greater than or equal to 160 dB re 1 μ Pa (rms) (for seismic airgun operations), and/or 180 dB re 1 μ Pa (rms) for cetaceans and 190 dB re 1 μ Pa (rms) for pinnipeds, with a discussion of the nature of the probable consequences of that exposure on the individuals that have been exposed.

(iv) A description of the implementation and effectiveness of the: (A) Terms and Conditions of the Biological Opinion's Incidental Take Statement (ITS) (attached); and (B) mitigation measures of the IHA. For the Biological Opinion, the report shall confirm the implementation of each Term and Condition, as well as any conservation recommendations, and describe their effectiveness, for minimizing the adverse effects of the action on Endangered Species Act-listed marine mammals.

(b) Submit a final report to the Chief, Permits and Conservation Division, Office of Protected Resources, NMFS, within 30 days after receiving comments from NMFS on the draft report. If NMFS decides that the draft report needs no comments, the draft report shall be considered to be the final report.

8. Reporting Prohibited Take

(a) (i) In the unanticipated event that the specified activity clearly causes the take of a marine mammal in a manner prohibited by this Authorization, such as an injury (Level A harassment), serious injury or mortality (*e.g.*, through ship-strike, gear interaction, and/or entanglement), SIO shall immediately cease the specified activities and immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401 and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the following information:

(ii) Time, date, and location (latitude/longitude) of the incident; the name and

type of vessel involved; the vessel's speed during and leading up to the incident; description of the incident; status of all sound source use in the 24 hours preceding the incident; water depth; environmental conditions (*e.g.*, wind speed and direction, Beaufort sea state, cloud cover, and visibility); description of marine mammal observations in the 24 hours preceding the incident; species identification or description of the animal(s) involved; the fate of the animal(s); and photographs or video footage of the animal (if equipment is available).

Activities shall not resume until NMFS is able to review the circumstances of the prohibited take. NMFS shall work with SIO to determine what is necessary to minimize the likelihood of further prohibited take and ensure MMPA compliance. SIO may not resume their activities until notified by NMFS via letter, email, or telephone.

Reporting an Injured or Dead Marine Mammal With an Unknown Cause of Death

(b) In the event that SIO discover an injured or dead marine mammal, and the lead PSO determines that the cause of the injury or death is unknown and the death is relatively recent (*i.e.*, in less than a moderate state of decomposition), SIO shall immediately report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov. The report must include the same information identified in Condition 8(c)(i) above. Activities may continue while NMFS reviews the circumstances of the incident. NMFS shall work with SIO to determine whether modifications in the activities are appropriate.

Reporting an Injured or Dead Marine Mammal Not Related to the Activities

(c) In the event that SIO discovers an injured or dead marine mammal, and the lead PSO determines that the injury or death is not associated with or related to the activities authorized in Condition 2 of this Authorization (*e.g.*, previously wounded animal, carcass with moderate to advanced decomposition, or scavenger damage), SIO shall report the incident to the Chief of the Permits and Conservation Division, Office of Protected Resources, NMFS, at 301-427-8401, and/or by email to Jolie.Harrison@noaa.gov and Howard.Goldstein@noaa.gov, within 24 hours of the discovery. SIO shall provide photographs or video footage (if available) or other documentation of the

stranded animal sighting to NMFS. Activities may continue while NMFS reviews the circumstances of the incident.

9. *Endangered Species Act Biological Opinion and Incidental Take Statement*

(a) SIO is required to comply with the Terms and Conditions of the ITS corresponding to NMFS's Biological Opinion issued to both NSF and SIO, and NMFS's Office of Protected Resources.

(b) A copy of this Authorization and the ITS must be in the possession of all

contractors and PSO(s) operating under the authority of this Incidental Harassment Authorization.

Request for Public Comments

NMFS requests comment on our analysis, the draft authorization, and any other aspect of the notice of the proposed IHA for SIO's low-energy seismic survey. Please include with your comments any supporting data or literature citations to help inform our final decision on SIO's request for an MMPA authorization. Concurrent with

the publication of this notice in the **FEDERAL REGISTER**, NMFS is forwarding copies of this application to the Marine Mammal Commission and its Committee of Scientific Advisors.

Dated: March 12, 2015.

Perry Gayaldo,

Deputy Director, Office of Protected Resources, National Marine Fisheries Service.

[FR Doc. 2015-06261 Filed 3-19-15; 8:45 am]

BILLING CODE 3510-22-P